

RATIONAL SOLUTIONS OF THE SCHLESINGER SYSTEM AND ISOPRINCIPAL DEFORMATIONS OF RATIONAL MATRIX FUNCTIONS I.

VICTOR KATSNELSON AND DAN VOLOK

ABSTRACT. In the present paper we discuss the general facts concerning the Schlesinger system: the τ -function, the local factorization of solutions of Fuchsian equations and holomorphic deformations. We introduce the terminology "isoprincipal" for the deformations of Fuchsian equations with general (not necessarily non-resonant) matrix coefficients corresponding to solutions of the Schlesinger system. Every isoprincipal deformation is isomonodromic. The converse is also true in the non-resonant case, but not in general.

In the forthcoming sequel we shall give an explicit description of a class of rational solutions of the Schlesinger system, based on the techniques developed here, and the realization theory for rational matrix functions.

NOTATIONS.

- $\mathbb{C}$ stands for the complex plane;
- $\overline{\mathbb{C}}$ stands for the extended complex plane (= the Riemann sphere):
 $\overline{\mathbb{C}} = \mathbb{C} \cup \infty$;
- $\mathbb{C}^n$ stands for the n - dimensional complex space;
- in the coordinate notation, a point $\mathbf{t} \in \mathbb{C}^n$ is written as $\mathbf{t} = (t_1, \dots, t_n)$;
- $\mathbb{C}_*^n$ is the set of those points $\mathbf{t} = (t_1, \dots, t_n) \in \mathbb{C}^n$, whose coordinates $t_1, \dots, t_n$ are pairwise different: $\mathbb{C}_*^n = \mathbb{C}^n \setminus (\bigcup_{1 \leq i, j \leq n, i \neq j} \{\mathbf{t} : t_i = t_j\})$;
- $\mathfrak{M}_k$ stands for the set of all $k \times k$ matrices with complex entries;
- $[\cdot, \cdot]$ denotes the commutator: for $A, B \in \mathfrak{M}_k$, $[A, B] = AB - BA$;
- I stands for the unity matrix of the appropriate dimension;
- $\mathcal{R}(\mathfrak{M}_k)$ stands for the set of all rational $\mathfrak{M}_k$ -valued functions R such that $\det R(z) \neq 0$;
- $\mathcal{P}(R)$ stands for the set of all poles of the function R , $\mathcal{N}(R)$ stands for the set of all poles of the function R^{-1} ; $\mathcal{P}(R)$ is said to be the pole set of the function R , $\mathcal{N}(R)$ is said to be the zero set of the function R .

Date: April 1, 2003.

1991 *Mathematics Subject Classification.* 34Mxx, 34M55, 93B15, 47A56.

Key words and phrases. Differential equations in the complex domain, monodromy, Schlesinger system.

The research of Victor Katsnelson was supported by the Minerva Foundation.

1. THE SCHLESINGER SYSTEM

The Schlesinger system is a Pfaffian system of differential equations of the form:

$$(1.1) \quad \begin{cases} \frac{\partial Q_i}{\partial t_j} = \frac{[Q_i, Q_j]}{t_i - t_j}, & 1 \leq i, j \leq n, \ i \neq j, \\ \frac{\partial Q_i}{\partial t_i} = - \sum_{\substack{1 \leq j \leq n \\ j \neq i}} \frac{[Q_i, Q_j]}{t_i - t_j}, & 1 \leq i \leq n, \end{cases} \quad \begin{matrix} (1.1.a) \\ (1.1.b) \end{matrix}$$

where $Q_1, \dots, Q_n$ are square $k \times k$ matrix functions of $\mathbf{t} = (t_1, \dots, t_n)$, $\mathbf{t} \in \mathbb{C}_*^n$. The system of equations (1.1) can be rewritten in a compact way as

$$(1.2) \quad \mathfrak{d}Q_\nu = \sum_{\substack{\mu=1 \\ \mu \neq \nu}}^n [Q_\mu, Q_\nu] \mathfrak{d} \log(t_\mu - t_\nu), \quad \nu = 1, \dots, n,$$

where $\mathfrak{d}$ is the exterior differential. The Schlesinger system (1.1) is over-determined. The compatibility (the integrability) condition for this system is of the form

$$(1.3) \quad \mathfrak{d} \left(\sum_{\substack{\mu=1 \\ \mu \neq \nu}}^n [Q_\mu, Q_\nu] \mathfrak{d} \log(t_\mu - t_\nu) \right) = 0, \quad \nu = 1, \dots, n.$$

The condition (1.3) must be fulfilled if $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ satisfy the equations (1.2). This condition expresses the equality $\mathfrak{d}(\mathfrak{d}Q_\nu) = 0$, which must be satisfied since $\mathfrak{d} \cdot \mathfrak{d} = 0$.

Let us present the integrability condition (1.3) in a more explicit form. Since $\mathfrak{d}\mathfrak{d} \log(t_\mu - t_\nu) = 0$,

$$(1.4) \quad \mathfrak{d} \left(\sum_{\substack{\mu=1 \\ \mu \neq \nu}}^n [Q_\mu, Q_\nu] \mathfrak{d} \log(t_\mu - t_\nu) \right) = \\ = \sum_{\substack{\mu=1 \\ \mu \neq \nu}}^n [\mathfrak{d}Q_\mu, Q_\nu] \wedge \mathfrak{d} \log(t_\mu - t_\nu) + \sum_{\substack{\mu=1 \\ \mu \neq \nu}}^n [Q_\mu, \mathfrak{d}Q_\nu] \wedge \mathfrak{d} \log(t_\mu - t_\nu),$$

where we have to substitute the expression of the form (1.2) for $\mathfrak{d}Q_\mu$ and $\mathfrak{d}Q_\nu$ in the right hand side of (1.4). Performing the substitutions, we obtain the expressions

$$(1.5) \quad \sum_{\substack{\mu=1 \\ \mu \neq \nu}}^n \left[\sum_{\substack{\lambda=1 \\ \lambda \neq \mu}}^n [Q_\lambda, Q_\mu], Q_\nu \right] \mathfrak{d} \log(t_\lambda - t_\mu) \wedge \mathfrak{d} \log(t_\mu - t_\nu) + \\ + \sum_{\substack{\mu=1 \\ \mu \neq \nu}}^n \left[Q_\mu, \sum_{\substack{\lambda=1 \\ \lambda \neq \nu}}^n [Q_\lambda, Q_\nu] \right] \mathfrak{d} \log(t_\lambda - t_\nu) \wedge \mathfrak{d} \log(t_\mu - t_\nu).$$

Since $\omega \wedge \omega = 0$ for every differential form ω , $\mathfrak{d} \log(t_\mu - t_\nu) \wedge \mathfrak{d} \log(t_\mu - t_\nu) = 0$. Therefore, the integrability condition for the Schlesinger system (1.2) can be

presented in the form

$$(1.6) \quad \sum_{\substack{\lambda=1, \mu=1; \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n [[Q_\lambda, Q_\mu], Q_\nu] \mathfrak{d} \log(t_\lambda - t_\mu) \wedge \mathfrak{d} \log(t_\mu - t_\nu) + \\ + \sum_{\substack{\lambda=1, \mu=1; \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n [Q_\mu, [Q_\lambda, Q_\nu]] \mathfrak{d} \log(t_\lambda - t_\nu) \wedge \mathfrak{d} \log(t_\mu - t_\nu) = 0, \quad \nu = 1, \dots, n.$$

Lemma 1.1. *For every holomorphic square matrix functions $Q_1, \dots, Q_n$, the integrability condition (1.6) holds.*

The proof of this lemma is based on two identities. The first of them is the Jakobi identity

$$(1.7) \quad [[A, B], C] + [[B, C], A] + [[C, A], B] = 0,$$

which holds for arbitrary square matrices A, B, C (of the same dimension). The second identity is an identity related to the differential holomorphic 1-form $\omega_{\lambda, \mu}$:

$$(1.8) \quad \omega_{\lambda, \mu} = \frac{dt_\lambda - dt_\mu}{t_\lambda - t_\mu}, \quad 1 \leq \lambda, \mu \leq n, \lambda \neq \mu.$$

Since

$$(1.9) \quad \omega_{\lambda, \mu} = \omega_{\mu, \lambda}, \quad 1 \leq \lambda, \mu \leq n, \lambda \neq \mu,$$

there are $\frac{n(n-1)}{2}$ different forms $\omega_{\lambda, \mu}$. The second identity is:

$$(1.10) \quad \omega_{\lambda, \mu} \wedge \omega_{\mu, \nu} + \omega_{\mu, \nu} \wedge \omega_{\nu, \lambda} + \omega_{\nu, \lambda} \wedge \omega_{\lambda, \mu} = 0, \quad 1 \leq \lambda, \mu, \nu \leq n, \quad \lambda \neq \mu \neq \nu.$$

To prove this identity we observe that the left-hand side of (1.10) is the sum of the expressions that are obtained from $\omega_{\lambda, \mu} \wedge \omega_{\mu, \nu}$ by all cyclic permutations of the indices λ, μ, ν . Computing the exterior product $\omega_{\lambda, \mu} \wedge \omega_{\mu, \nu}$:

$$(1.11) \quad \omega_{\lambda, \mu} \wedge \omega_{\mu, \nu} = \frac{dt_\lambda \wedge dt_\mu + dt_\mu \wedge dt_\nu + dt_\nu \wedge dt_\lambda}{(t_\lambda - t_\mu)(t_\mu - t_\nu)},$$

we see that it is represented as a fraction whose numerator is invariant with respect to cyclic permutations of the indices λ, μ, ν . Therefore, the right-hand side of (1.10) is equal to

$$\left\{ \frac{1}{(t_\lambda - t_\mu)(t_\mu - t_\nu)} + \frac{1}{(t_\mu - t_\nu)(t_\nu - t_\lambda)} + \frac{1}{(t_\nu - t_\lambda)(t_\lambda - t_\mu)} \right\} \\ \times (dt_\lambda \wedge dt_\mu + dt_\mu \wedge dt_\nu + dt_\nu \wedge dt_\lambda)$$

Evidently, the expression in the curly brackets vanishes. The identity (1.10) is proved.

Remark 1.2. The forms $\frac{1}{2\pi i} \omega_{\lambda, \mu}$, $1 \leq \lambda, \mu \leq n$, $\lambda \neq \mu$, generate the cohomology ring $H^*(\mathbb{C}_*^n, \mathbb{Z})$ of the space $\mathbb{C}_*^n$. This result was established by V. Arnold, [Arn].

PROOF of LEMMA 1.1. Let us fix $\nu \in 1, \dots, n$. First, we transform the first summand in (1.6). Applying the Jacobi identity to the factor $[[Q_\lambda, Q_\mu], Q_\nu]$ in the first sum, we rewrite (1.6) as

$$\begin{aligned}
 (1.12) \quad & \sum_{\substack{\lambda=1, \mu=1; \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n [[Q_\mu, Q_\nu], Q_\lambda] \omega_{\lambda, \mu} \wedge \omega_{\mu, \nu} \\
 & + \sum_{\substack{\lambda=1, \mu=1; \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n [[Q_\nu, Q_\lambda], Q_\mu] \omega_{\lambda, \mu} \wedge \omega_{\mu, \nu} \\
 & + \sum_{\substack{\lambda=1, \mu=1; \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n [[Q_\lambda, Q_\nu], Q_\mu] \omega_{\lambda, \nu} \wedge \omega_{\mu, \nu} = 0
 \end{aligned}$$

Since the summation runs over all pairs λ, μ , we can interchange λ and μ in the second and third sums of (1.12). Using also (1.9) and the anti-commutativity of the exterior product ($\omega \wedge \sigma = -\sigma \wedge \omega$ for every differential 1-forms ω and σ), we rewrite (1.12) as

$$\begin{aligned}
 (1.13) \quad & \sum_{\substack{\lambda=1, \mu=1; \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n [[Q_\mu, Q_\nu], Q_\lambda] \omega_{\lambda, \mu} \wedge \omega_{\mu, \nu} \\
 & + \sum_{\substack{\lambda=1, \mu=1; \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n [[Q_\mu, Q_\nu], Q_\lambda] \omega_{\nu, \lambda} \wedge \omega_{\lambda, \mu} \\
 & + \sum_{\substack{\lambda=1, \mu=1; \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n [[Q_\mu, Q_\nu], Q_\lambda] \omega_{\mu, \nu} \wedge \omega_{\nu, \lambda} = 0
 \end{aligned}$$

The last equality holds because its left-hand side is of the form

$$\sum_{\substack{\lambda=1, \mu=1; \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n [[Q_\mu, Q_\nu], Q_\lambda] \{ \omega_{\lambda, \mu} \wedge \omega_{\mu, \nu} + \omega_{\nu, \lambda} \wedge \omega_{\lambda, \mu} + \omega_{\mu, \nu} \wedge \omega_{\nu, \lambda} \},$$

where the sums in the curly brackets vanish according to the identity (1.10).

Q.E.D.

From Lemma 1.1 and from Frobenius theorem it follows that the Cauchy problem for the Schlesinger system is locally solvable for every initial data. More precisely, the following result holds:

Theorem 1.3. *Let $\mathbf{t}^0 \in \mathbb{C}_*^n$, $Q_1^0, \dots, Q_n^0 \in \mathfrak{M}_k$. Then there exist a neighborhood $\mathcal{V}(\mathbf{t}^0, \delta)$ of the point $\mathbf{t}^0$,*

$$\mathcal{V}(\mathbf{t}^0, \delta) = \{ \mathbf{t} \in \mathbb{C}_n : |t_1 - t_1^0| < \delta, \dots, |t_n - t_n^0| < \delta \}, \mathcal{V}(\mathbf{t}^0, \delta) \subset \mathbb{C}_*^n,$$

and matrix functions $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$,

$$Q_\nu(\mathbf{t}) : \mathcal{V}(\mathbf{t}^0, \delta) \mapsto \mathfrak{M}_k, \nu = 1, \dots, n,$$

which are holomorphic in $\mathcal{V}(\mathbf{t}^0, \delta)$, satisfy the Schlesinger system (1.1) for $\mathbf{t} \in \mathcal{V}(\mathbf{t}^0, \delta)$, and meet the initial condition

$$(1.14) \quad Q_\nu(\mathbf{t}^0) = Q_\nu^0, \nu = 1, \dots, n$$

at the point $\mathbf{t}^0$. Here δ is a positive number which depends on $\mathbf{t}^0$ and $Q_\nu^0, \nu = 1, \dots, n$.

Thus Theorem 1.3 is a local existence result. Actually, a much more stronger global existence result holds for the Schlesinger system. It turns out that the solution $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ of the above introduced Cauchy problem for this system exists globally as meromorphic (multi-valued) functions in $\mathbb{C}_*^n$. We do not discuss this more general result in detail – this will lead us too far. However, it is worth mentioning that every solution of the Schlesinger system can be represented as a fraction of holomorphic functions in $\mathbb{C}_*^n$, where some complex valued holomorphic function $\tau(\cdot)$ on $\mathbb{C}_*^n$ appears in the denominator. This is the so-called τ -function for the considered solution. The singularities of this solution are precisely those $\mathbf{t} \in \mathbb{C}_*^n$ where $\tau(\mathbf{t}) = 0$. The "zeros" of the function $\tau(\cdot)$ are responsible for the singularities of the solution $Q_1(\cdot), \dots, Q_n(\cdot)$.

2. THE TAU-FUNCTION OF THE SCHLESINGER SYSTEM

Lemma 2.1. *Let matrix-functions $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ satisfy the Schlesinger system (1.1). Then the differential 1-form ω :*

$$(2.1) \quad \omega \stackrel{\text{def}}{=} \frac{1}{2} \sum_{\substack{\lambda, \mu=1 \\ \lambda \neq \mu}}^n \text{tr}(Q_\lambda(\mathbf{t})Q_\mu(\mathbf{t})) \mathfrak{d} \log(t_\lambda - t_\mu),$$

is closed:

$$(2.2) \quad \mathfrak{d}\omega = 0.$$

PROOF. It is clear that

$$\begin{aligned} \mathfrak{d}\omega &= \frac{1}{2} \sum_{\substack{\lambda, \mu=1 \\ \lambda \neq \mu}}^n \text{tr} \left((\mathfrak{d}Q_\lambda(t)) Q_\mu(t) \right) \mathfrak{d} \log(t_\lambda - t_\mu) \\ &\quad + \frac{1}{2} \sum_{\substack{\lambda, \mu=1 \\ \lambda \neq \mu}}^n \text{tr} \left(Q_\lambda(t) (\mathfrak{d}Q_\mu(t)) \right) \mathfrak{d} \log(t_\lambda - t_\mu). \end{aligned}$$

Interchanging the summation indices λ and μ in the first sum (and using the rule $\text{tr}AB = \text{tr}BA$), we obtain (after renaming $\lambda \rightarrow \nu, \nu \rightarrow \lambda$) that

$$(2.3) \quad \mathfrak{d}\omega = \sum_{\substack{\lambda, \nu=1 \\ \lambda \neq \nu}}^n \text{tr} \left(Q_\lambda(t) (\mathfrak{d}Q_\nu(t)) \right) \mathfrak{d} \log(t_\nu - t_\lambda).$$

Substituting the expression (1.2) for $\mathfrak{d}Q_\nu$ into (2.3), we obtain the following expression for $\mathfrak{d}\omega$:

$$(2.4) \quad \mathfrak{d}\omega = \sum_{\substack{\lambda=1, \mu=1, \nu=1 \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n \text{tr} \left(Q_\lambda [Q_\mu, Q_\nu] \right) \omega_{\mu, \nu} \wedge \omega_{\nu, \lambda},$$

where $\omega_{\lambda,\mu}$ is defined by (1.8), or

$$(2.5) \quad \mathfrak{d}\omega = \sum_{\substack{\lambda=1, \mu=1, \nu=1 \\ \lambda \neq \mu, \lambda \neq \nu, \mu \neq \nu}}^n \text{tr} \left(Q_\lambda Q_\mu Q_\nu - Q_\lambda Q_\nu Q_\mu \right) \omega_{\mu,\nu} \wedge \omega_{\nu,\lambda}.$$

The traces in the last formula are invariant under a circular permutation of the three indices λ, μ, ν . So we can replace $\omega_{\mu,\nu} \wedge \omega_{\nu,\lambda}$ by $\frac{1}{3}(\omega_{\mu,\nu} \wedge \omega_{\nu,\lambda} + \omega_{\nu,\lambda} \wedge \omega_{\lambda,\mu} + \omega_{\lambda,\mu} \wedge \omega_{\mu,\nu})$. But this quantity vanishes identically according to the identity (1.10). Thus, $\mathfrak{d}\omega = 0$. Q.E.D.

Since a closed form $\mathfrak{d}\omega$ is locally exact, the following theorem is an immediate consequence of Lemma 2.1:

Theorem 2.2. *Let holomorphic matrix-functions $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ satisfy the Schlesinger system (1.1) in some simply connected domain $\mathcal{D}, \mathcal{D} \subset \mathbb{C}_*^n$. Then there exists a holomorphic non-vanishing function $\tau(\mathbf{t})$ in $\mathcal{D}$ such that*

$$(2.6) \quad \mathfrak{d} \log \tau(\mathbf{t}) = \frac{1}{2} \sum_{\substack{\lambda, \mu=1 \\ \lambda \neq \mu}}^n \text{tr}(Q_\lambda(\mathbf{t}) Q_\mu(\mathbf{t})) \mathfrak{d} \log(t_\lambda - t_\mu)$$

Such a function τ is unique up to a constant factor: if $\tau_1(\mathbf{t})$ and $\tau_2(\mathbf{t})$ are two functions satisfying the relation (2.6), then $\tau_2(\mathbf{t}) = c\tau_1(\mathbf{t})$, where $c \neq 0$ is a constant.

Definition 2.3. A function τ related to a solution $Q_1, \dots, Q_n$ of the Schlesinger system (1.1) by the equality (2.6) is said to be the τ -function for this solution.

The exactness of the differential form (2.1), stated in our Lemma 2.1, was established by M. Sato, T. Miwa and M. Jimbo in [SMJ2] (see §2.4 there). In [SMJ2], the potential (2.6) was introduced as well as the term " τ -function" for this potential.

Remark 2.4. From Lemma 2.1 it follows that the τ -function for a solution of the Schlesinger equation is a holomorphic non-vanishing (multi-valued) function for those $\mathbf{t}$ where the solution is holomorphic. Actually, the τ -function is holomorphic on the whole $\mathbb{C}_*^n$. It vanishes for those $\mathbf{t}$ where the solution is singular. (See the last paragraph of the section 1). In a very general setting, the holomorphy of τ -functions was proved in [Miw].

Remark 2.5. The τ -function is a very general concept related to non-linear differential equations. This function was first discovered by R. Hirota as a tool for generating many-soliton solutions. (References to the works of H. Hirota can be found in bibliography of the book [New]). Let us give examples of τ -functions for the Korteweg - deVries equation

$$(2.7) \quad u_t - 6uu_x + u_{xxx} = 0.$$

N - soliton solution

$$(2.8) \quad u(x, t) = -2 \frac{\partial^2}{\partial x^2} \log \Delta(x, t),$$

$$(2.9) \quad \Delta(x, t) = \det \left(\delta_{jk} + \frac{c_j c_k}{\eta_j + \eta_k} \exp[-(\eta_j^3 + \eta_k^3)t - (\eta_j + \eta_k)x] \right)_{j,k=1,\dots,N};$$

N - phase solution

$$(2.10) \quad u(x, t) = -2 \frac{\partial^2}{\partial x^2} \log \vartheta(\mathbf{a}x + \mathbf{b}t + \mathbf{c}),$$

where ϑ is the Riemann theta function;
rapidly decreasing solution

$$(2.11) \quad u(x, t) = -2 \frac{\partial^2}{\partial x^2} \log \Delta_{[x, \infty)}(t),$$

with $\Delta_{[x, \infty)}(t)$ the Fredholm determinant of the Gelfand - Levitan equation. In these examples the τ -function is $\Delta(x, t)$, $\vartheta(\mathbf{a}x + \mathbf{b}t + \mathbf{c})$, $\Delta_{[x, \infty)}(t)$, respectively.

In the paper [WNTB], τ -function appeared as the spin - spin correlation function for the two - dimensional Ising model. In [JMMS], the Fredholm determinant $\det(\mathbf{I} - \lambda \mathbf{K})$ of the integral operator $\mathbf{K}$ with the kernel $K(x, x') = \frac{\sin(x-x')}{x-x'}$ on the union of intervals $\cup_{1 \leq j \leq n} [a_{2j-1}, a_{2j}]$, $a_1 < \dots < a_{2n}$, is expressed as the τ -function $\tau(a_1, \dots, a_{2n})$ of the Schlesinger system constructed from the operator. (See in particular section 7 of [JMMS], where the deformation theory of the Fredholm integral operator with the kernel $\frac{\sin(x-x')}{x-x'}$ was related to the deformation theory of linear differential equations). The interrelation between the deformation theory of linear differential operators, the Schlesinger system and its τ -function, correlation function in statistical physics theory, quantum fields and Fredholm determinants are covered in the expository survey [SMJ1]. The τ -function for general matrix linear differential equation with rational coefficients was considered in [JMU], and was studied in great detail in [JM1], [JM2].

However, the importance of the τ -function and its central role in the soliton theory was not understood before the works of the research group from Kyoto (M. Sato, T. Miwa, M. Jimbo, E. Date, M. Kashiwara, Y. Sato). The work of this group, surrounding Mikio Sato, reached its peak in the years around 1981. In the paper [SaSa], the time evolution of a solution of a non-linear equation is interpreted as the dynamical motion of a point of the infinite dimensional Grassman manifold. In the papers of E. Date, M. Jimbo, M. Kashiwara and T. Miwa, Lie algebras and vertex operators were incorporated into the soliton theory. (See the series of seven notes [DJKM1], and the paper [DJKM2]). In this setting, the τ -function appears in a very general way. The paper [Sa] explains some ideas from [SaSa] in more detail. The further development of the approach related to the consideration of soliton equations as dynamic systems on an infinite dimensional Grassman manifold is done in [SeWi]. In particular, in section 3 of [SeWi] the geometric meaning of the τ -function is explained: this function appears as an infinite determinant related to the determinant bundle over the Grassman manifold. In [MSW], the τ -function is defined as the generalized cross-ratio of four points of the Grassman manifold. Tau function is discussed in Chapters 7 and 8 of the book [Dick]. The book [MJD] is a presentation of the algebraic analysis of the nonlinear differential equations of the KdV type written for students. In particular, it contains a very clear presentation of the circle of problem related to τ -functions. Chapter 4 of the book [New] is dedicated to the τ -function and its relation to the Painlevé property and to the Bäcklund transformations. This non-formally written book covers a very widespread circle of problems.

In the present work we construct a class of rational solutions of the Schlesinger system (1.1). For such a solution, we calculate its τ -function explicitly. This τ -function is equal to the determinant of some (finite) matrix which is intimately related to this solution. In the considered special case, τ -function is a rational function of the variables $t_1, \dots, t_n$.

3. THE POTENTIAL FUNCTION FOR A SOLUTION OF THE SCHLESINGER SYSTEM.

From (1.1.b) it follows that if the matrix functions $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ satisfy the Schlesinger system in a domain $\mathcal{G} \in \mathbb{C}_*^n$, then

$$(3.1) \quad \frac{\partial Q_i(\mathbf{t})}{\partial t_j} \equiv \frac{\partial Q_j(\mathbf{t})}{\partial t_i}.$$

Hence, there exists a matrix function $V(\mathbf{t})$ in $\mathcal{G}$ such that

$$(3.2) \quad Q_j(\mathbf{t}) = \frac{\partial V(\mathbf{t})}{\partial t_j}, \quad \mathbf{t} \in \mathcal{G}, \quad j = 1, \dots, n.$$

Remark 3.1. If the domain $\mathcal{G}$ is multiconnected, the function $V(\mathbf{t})$ can be multivalued. In this case we can consider $V(\mathbf{t})$ as single-valued function on the universal covering $\text{cov}(\mathcal{G}, \mathbf{t}_0)$ of the domain $\mathcal{G}$, where $\mathbf{t}_0$ is a chosen distinguished point of $\mathcal{G}$.

Definition 3.2. The matrix function $V(\mathbf{t})$ which satisfies the condition (3.2) is said to be the *potential matrix function* for the solution $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ of the Schlesinger system.

We have proved that the potential matrix function exists for any solution of the Schlesinger system. It is clear that the potential function is unique up to an arbitrary constant summand.

Lemma 3.3. *Let matrices $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ satisfy the Schlesinger system (1.1), and moreover the condition*

$$(3.3) \quad \sum_{1 \leq i \leq n} Q_i(\mathbf{t}) = 0$$

for every $\mathbf{t} \in \mathcal{G}$.

Then the function

$$(3.4) \quad V(\mathbf{t}) \stackrel{\text{def}}{=} \sum_{1 \leq i \leq n} t_i \cdot Q_i(\mathbf{t}), \quad \mathbf{t} \in \mathcal{G},$$

is a potential matrix function for the solution $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$.

PROOF. Differentiating the expression on the left hand side of (3.4), we obtain

$$(3.5) \quad \frac{\partial V(\mathbf{t})}{\partial t_j} = Q_j(\mathbf{t}) + \sum_{1 \leq i \leq n} t_i \cdot \frac{\partial Q_i(\mathbf{t})}{\partial t_j}.$$

Thus, we have to prove that

$$(3.6) \quad \sum_{1 \leq i \leq n} t_i \cdot \frac{\partial Q_i(\mathbf{t})}{\partial t_j} = 0.$$

Substituting the expression (1.1) for $\frac{\partial Q_i(\mathbf{t})}{\partial t_j}$ into the left hand side of (3.6), we obtain

$$\begin{aligned}
 (3.7) \quad \sum_{1 \leq i \leq n} t_i \cdot \frac{\partial Q_i(\mathbf{t})}{\partial t_j} &= \sum_{\substack{1 \leq i \leq n, \\ i \neq j}} t_i \cdot \frac{[Q_i(\mathbf{t}), Q_j(\mathbf{t})]}{t_i - t_j} - t_j \cdot \sum_{\substack{1 \leq i \leq n, \\ i \neq j}} \frac{[Q_i(\mathbf{t}), Q_j(\mathbf{t})]}{t_i - t_j} \\
 &= \sum_{\substack{1 \leq i \leq n, \\ i \neq j}} [Q_i(\mathbf{t}), Q_j(\mathbf{t})] = \sum_{1 \leq i \leq n} [Q_i(\mathbf{t}), Q_j(\mathbf{t})] = 0.
 \end{aligned}$$

Thus, (3.6) holds.

Q.E.D.

Lemma 3.4. *Let $\mathcal{G}$ be a domain in $\mathbb{C}_*^n$, and let matrix functions $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ satisfy the Schlesinger system (1.1) for $\mathbf{t} \in \mathcal{G}$. Assume that the equality (3.3) holds at least at one point $\mathbf{t} = \mathbf{t}_0 \in \mathcal{G}$. Then this equality holds at every point $\mathbf{t} \in \mathcal{G}$.*

PROOF. The equality (3.3) will be proved if we prove that

$$(3.8) \quad \frac{\partial}{\partial t_j} \left(\sum_{1 \leq i \leq n} Q_i(\mathbf{t}) \right) = 0 \quad \text{for } \mathbf{t} \in \mathcal{G}.$$

From (1.1) it follows that

$$\begin{aligned}
 \sum_{1 \leq i \leq n} \frac{\partial Q_i(\mathbf{t})}{\partial t_j} &= \frac{\partial Q_j(\mathbf{t})}{\partial t_j} + \sum_{\substack{1 \leq i \leq n, \\ i \neq j}} \frac{\partial Q_i(\mathbf{t})}{\partial t_j} \\
 &= - \sum_{\substack{1 \leq i \leq n, \\ i \neq j}} \frac{[Q_i(\mathbf{t}), Q_j(\mathbf{t})]}{t_i - t_j} + \sum_{\substack{1 \leq i \leq n, \\ i \neq j}} \frac{[Q_i(\mathbf{t}), Q_j(\mathbf{t})]}{t_i - t_j} = 0 \quad \text{for } \mathbf{t} \in \mathcal{G}.
 \end{aligned}$$

Thus, (3.8) holds.

Q.E.D.

The following lemma is an immediate consequence of two previous ones.

Lemma 3.5. *Let the matrix functions $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ satisfy the Schlesinger system (1.1) for every $\mathbf{t} \in \mathcal{G}$, where $\mathcal{G}$ is a domain in $\mathbb{C}_*^n$. Assume that the condition (3.3) holds at least for one point $\mathbf{t} = \mathbf{t}_0 \in \mathcal{G}$.*

Then the function $V(\mathbf{t})$ which is defined by (3.4) is a potential matrix function for the solution $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$.

The expression $\sum t_i Q_i(\mathbf{t})$, which appears on the right hand side of the equality (3.4), has a useful interpretation in terms of an auxiliary ordinary differential equation. Let $\mathbf{t} = (t_1, \dots, t_n) \in \mathbb{C}_*^n$, and $Q_1, \dots, Q_n$ be $k \times k$ matrices. (For the moment, we consider the point $\mathbf{t}$ and the matrices $Q_1, \dots, Q_n$ as fixed rather than variable.) Let us introduce an auxiliary complex variable x belonging to the domain ${}^1\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$. Consider the

¹We recall that $\overline{\mathbb{C}} = \mathbb{C} \cup \infty$ is the extended complex plane.

linear differential equation

$$(3.9) \quad \frac{dY(x)}{dx} = \left(\sum_{1 \leq i \leq n} \frac{Q_i}{x - t_i} \right) \cdot Y(x)$$

with respect to a $k \times k$ matrix function $Y(x)$ of the variable x . The coefficient matrix function $\sum \frac{Q_i}{x - t_i}$ of the equation (3.9) is holomorphic in the domain $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, and if the condition

$$(3.10) \quad \sum_{1 \leq i \leq n} Q_i = 0$$

holds, then the residue of the coefficient matrix function at the point ∞ vanishes. Thus, under the condition (3.10), the Cauchy problem with the initial condition

$$(3.11) \quad Y(\infty) = I$$

for the differential equation (3.9) in the domain $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$ is solvable. The solution $Y(x)$ of this Cauchy problem is a function holomorphic in the disk $R < |x| \leq \infty$, where $R = \max_j |t_j|$. The function $Y(x)$ admits a Laurent expansion in this disk:

$$(3.12) \quad Y(x) = I + \sum_{1 \leq l \leq \infty} \frac{Y_{-l}}{x^l}, \quad R < |x| \leq \infty.$$

Lemma 3.6. *Under the condition (3.10), the Laurent coefficient Y_{-1} of the expansion (3.12) for the solution $Y(x)$ of the Cauchy problem (3.9) - (3.11) can be expressed as*

$$(3.13) \quad Y_{-1} = - \sum_{1 \leq i \leq n} t_i Q_i.$$

PROOF. Under the condition (3.10), the Laurent expansion at ∞ of the coefficient matrix function $\sum \frac{Q_i}{x - t_i}$ is of the form

$$(3.14) \quad \sum_{1 \leq i \leq n} \frac{Q_i}{x - t_i} = \left(\sum_{1 \leq i \leq n} t_i Q_i \right) \cdot x^{-2} + o(|x|^{-2}) \quad \text{as } x \rightarrow \infty.$$

Substituting the expansions (3.12) and (3.14) into the differential equation (3.9) and comparing the Laurent coefficients, we obtain (3.13).

Q.E.D.

Theorem 3.7. *Let $k \times k$ matrix functions $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ satisfy the Schlesinger system for $\mathbf{t} \in \mathcal{G}$, where $\mathcal{G}$ is a domain in $\mathbb{C}_*^n$. Let the equality (3.3) hold at least at one point $\mathbf{t} = \mathbf{t}_0 \in \mathcal{G}$, or, what is the same ², at every*

²Lemma 3.4

point $\mathbf{t} \in \mathcal{G}$. Assume that for each fixed $\mathbf{t} \in \mathcal{G}$, $k \times k$ matrix function $Y(x, \mathbf{t})$ satisfies the linear differential equation

$$(3.15) \quad \frac{dY(x, \mathbf{t})}{dx} = \left(\sum_{1 \leq i \leq n} \frac{Q_i(\mathbf{t})}{x - t_i} \right) \cdot Y(x, \mathbf{t}), \quad x \in \mathbb{C} \setminus \{t_1, \dots, t_n\},$$

with respect to x and the initial condition ³

$$(3.16) \quad Y(x, \mathbf{t})|_{x=\infty} = I.$$

Then the coefficient ⁴ $V(\mathbf{t})$ of the Laurent expansion

$$(3.17) \quad Y(x, \mathbf{t}) = I - \frac{V(\mathbf{t})}{x} + \frac{Y_{-2}(\mathbf{t})}{x^2} + \frac{Y_{-3}(\mathbf{t})}{x^3} + \dots, \quad (\max_i |t_i|) < |x| \leq \infty,$$

is the potential matrix function for the solution $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$, i.e. the equalities (3.2) hold.

PROOF. This theorem is an immediate consequence of Lemmas 3.3 and 3.6. Q.E.D.

The differential equation (3.15) plays an important role in what follows.

4. THE SCHLESINGER SYSTEM AS THE COMPATIBILITY CONDITION FOR SOME OVER-DETERMINED SYSTEM OF LINEAR DIFFERENTIAL EQUATIONS.

The result stated in this section is an important step in the study of the nonlinear Schlesinger system (1.1) by linear methods.

Theorem 4.1. *Let $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ be $k \times k$ matrix functions, holomorphic with respect to $\mathbf{t} = (t_1, \dots, t_n)$ in some open domain $\mathcal{G}$, $\mathcal{G} \subseteq \mathbb{C}_*^n$.*

Assume that the following system of linear differential equations

$$(4.1a) \quad \frac{\partial Y}{\partial x} = \left(\sum_{1 \leq j \leq n} \frac{Q_j(\mathbf{t})}{x - t_j} \right) Y,$$

$$(4.1b) \quad \frac{\partial Y}{\partial t_k} = -\frac{Q_k(\mathbf{t})}{x - t_k} Y \quad (k = 1, \dots, n)$$

for a square matrix function Y is compatible, that is, there exists a $k \times k$ matrix function $Y = Y(x; \mathbf{t})$, which is holomorphic, satisfies the equations (4.1) for $x \in \mathbb{C}$, $\mathbf{t} \in \mathcal{G}$, $x \neq t_1, \dots, t_n$, and, moreover, is not degenerate:

$$(4.2) \quad \det Y(x; \mathbf{t}) \neq 0, \quad \text{for } x \in \mathbb{C}, \mathbf{t} \in \mathcal{G}, x \neq t_1, \dots, x \neq t_n.$$

Then the matrix functions $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ satisfy the Schlesinger system (1.1) for $\mathbf{t} \in \mathcal{G}$.

³Due to (3.3), the value $Y(x, \mathbf{t})|_{x=\infty}$ exists.

⁴Here we redenote the Laurent coefficient $Y_{-1}(\mathbf{t})$ by $V(\mathbf{t})$.

PROOF. Let us exploit the equalities

$$(4.3) \quad \frac{\partial}{\partial t_k} \frac{\partial Y}{\partial x} = \frac{\partial}{\partial x} \frac{\partial Y}{\partial t_k}, \quad 1 \leq k \leq n.$$

Differentiating equation(4.1a) with respect to t_k , we obtain

$$\frac{\partial}{\partial t_k} \frac{\partial Y}{\partial x} = \left(\sum_{1 \leq j \leq n} \frac{Q_j}{x - t_j} \right) \frac{\partial Y}{\partial t_k} + \left(\sum_{1 \leq j \leq n} \frac{\frac{\partial Q_j}{\partial t_k}}{x - t_j} \right) Y + \frac{Q_k}{(x - t_k)^2} Y.$$

Substituting the expression for $\frac{\partial Y}{\partial t_k}$ from (4.1b) into the last equality, we get

$$(4.4) \quad \frac{\partial}{\partial t_k} \frac{\partial Y}{\partial x} = - \sum_{1 \leq j \leq n} \frac{Q_j Q_k}{(x - t_j)(x - t_k)} Y + \frac{Q_k}{(x - t_k)^2} Y + \sum_{1 \leq j \leq n} \frac{\frac{\partial Q_j}{\partial t_k}}{x - t_j} Y.$$

Differentiating equation(4.1b) with respect to x , we obtain

$$\frac{\partial}{\partial x} \frac{\partial Y}{\partial t_k} = \frac{Q_k}{(x - t_k)^2} Y - \frac{Q_k}{x - t_k} \frac{\partial Y}{\partial x}.$$

Substituting the expression for $\frac{\partial Y}{\partial x}$ from (4.1a) into the last equality, we get

$$(4.5) \quad \frac{\partial}{\partial x} \frac{\partial Y}{\partial t_k} = \frac{Q_k}{(x - t_k)^2} Y - \sum_{1 \leq j \leq n} \frac{Q_k Q_j}{(x - t_j)(x - t_k)} Y$$

Thus, equation (4.3) takes the form

$$(4.6) \quad \sum_{1 \leq l \leq n} \frac{\frac{\partial Q_l}{\partial t_k}}{x - t_l} = \sum_{\substack{1 \leq l \leq n \\ l \neq k}} \frac{[Q_l, Q_k]}{(x - t_l)(x - t_k)}$$

(Since matrix Y is non-degenerate, we can cancel out the factor Y .)

The numerators in (4.6) do not depend on x . Thus, (4.6) expresses the equality of two rational functions. Comparing the residues of both sides of (4.6) at $x = t_j, j \neq k$, we obtain (1.1.a). Comparing the residues at $x = t_k$, we obtain (1.1.b). Q.E.D.

Remark 4.2. In the proof of Theorem 4.1 we have not exploited the equalities

$$\frac{\partial}{\partial t_k} \frac{\partial Y}{\partial t_j} = \frac{\partial}{\partial t_j} \frac{\partial Y}{\partial t_k}, \quad 1 \leq k, j \leq n.$$

They lead to (1.1.a) and provide no additional information on $Q_1, \dots, Q_n$. Thus, reversing the reasoning in the proof, we can demonstrate that the Schlesinger system (1.1) is not only a necessary, but also a sufficient condition for the solvability of the linear system (4.1a) - (4.1b) for a non-degenerate matrix function Y .

Remark 4.3. The result, which we formulated as Theorem 4.1, is contained in the paper [Sch1] of L. Schlesinger. (See Section II of this paper, especially p. 294. See also p. 324 of the book [Sch2]). This was the paper [Sch1] where the Schlesinger equations (1.1) originally appeared. In [Sch1], the equations (1.1) were derived as the compatibility condition of some over-determined system of linear differential equations. In [Gar], R. Garnier has derived the Painlevé equation P_{VI} from the compatibility condition of the appropriate over-determined system of linear differential equations. (This over-determined system itself appeared in the paper [FuR] of R. Fuchs, which, in its turn, was based on works of his father L. Fuchs published in *Sitzungsberichte der Berliner Akademie der Wissenschaften* in 1888 - 1898). Since the publication of the works [ZaSh] and [AKNS], representing a solution of a nonlinear differential equation as the compatibility condition for some over-determined system of linear differential equations has become a common wisdom. In [AKNS], under the strong influence of the paper [ZaSh], M. Ablowitz, D. Kaup, A. Newell and H. Segur, have represented the Korteweg de Vries equation, the nonlinear Schrödinger equation, the Sine-Gordon equation and a number of other non-linear equations of physical significance as the compatibility conditions for the appropriate systems of linear differential equations.

5. FUCHSIAN EQUATIONS AND BEHAVIOR OF THEIR SOLUTIONS – GLOBAL AND LOCAL

A Fuchsian equation is a linear differential equation of the form

$$(5.1) \quad \frac{dY}{dx} = \left(\sum_{1 \leq j \leq n} \frac{Q_j}{x - t_j} \right) Y,$$

where $t_1, \dots, t_n$ are pairwise distinct points of the complex plane ($t_1 \neq t_2 \neq \dots \neq t_n$), $Q_j, j = 1, \dots, n$ are square matrices of the same dimension (say, $k \times k$), x is the complex variable which belongs to the punctured Riemann sphere $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$. The points $t_1, \dots, t_n$ are regular singular points for the equation (5.1). The residue of the coefficient matrix $\sum_{1 \leq k \leq n} \frac{Q_k}{x - t_k}$ of the equation (5.1) at the point ∞ is equal to $-\sum_{1 \leq k \leq n} Q_k$. So, if the condition

$$(5.2) \quad \sum_{1 \leq k \leq n} Q_k = 0$$

is satisfied, the point ∞ is a regular point for the equation (5.1). In what follows we assume that the condition (5.2) is satisfied.

We consider the differential equation (5.1) as a *matricial* one. This means that we consider solutions $Y(x)$ of this equation which are $k \times k$ matrix functions. If x_0 is a regular point for the equation (5.1) (that is $x_0 \neq t_1, \dots, t_n$), then for every $k \times k$ matrix Y_0 there exists the solution $Y(x; x_0, Y_0)$

which satisfies the initial condition

$$(5.3) \quad Y(x; x_0, Y_0)|_{x=x_0} = Y_0.$$

This solution is a holomorphic function in a disk neighborhood of the initial point x_0 . It can be continued analytically along any path which does not pass through the singular points $t_1, \dots, t_n$. Thus the solution of the *linear* differential equation (5.1) exists *globally* and is a (multi-valued) holomorphic function on the punctured Riemann sphere $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$. For a point $x \in \overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, the value of the analytic continuation of the solution $Y(\cdot; x_0, Y_0)$ along a path $\gamma(x_0, x) \subset \overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, which starts at the point x_0 and ends at the point x , can depend on this path γ . If $\gamma'(x_0, x)$ and $\gamma''(x_0, x)$ are two such paths, homotopic in $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, then the values of analytic continuations of the solution $Y(\cdot; x_0, Y_0)$ along γ' and γ'' coincide. Thus, the value of the analytic continuation of the solution $Y(\cdot; x_0, Y_0)$ to the point x depends on *the homotopy class of the path* leading from x_0 to x , along which the continuation is made, but *not on the path itself*. In other words, the solution $Y(\cdot, x_0, Y_0)$ of the differential equation (5.1) with the initial condition (5.3) is holomorphic function on the universal covering space $\text{cov}(\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}; x_0)$ of the space $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$ with the distinguished point x_0 .

Definition 5.1. A solution $Y(x)$ of a linear differential equation in a connected domain is said to be **fundamental** if $\det Y(x) \not\equiv 0$ (or, what is equivalent, if $\det Y(x) \neq 0$ for some point x_0 from this domain).

If $Y(x)$ is a fundamental solution of a linear differential equation, and $Z(x)$ is another solution of this equation then $Z(x) = Y(x)C$, where C is a constant (with respect to x) matrix. In what follows we distinguish the solution $Y(x)$ of the differential equation (5.1), which satisfies the normalizing (initial) condition

$$(5.4) \quad Y(x_0) = I$$

at a distinguished point x_0 which is regular for the equation (5.1).

Let $\gamma(x_0, x_0)$ be a *closed* path, which starts at the point x_0 and ends at *the same point* x_0 . We call such a path *a loop with the distinguished point* x_0 . Let $x_0^*(\gamma)$ be the point of the universal covering space $\text{cov}(\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}; x_0)$, which corresponds to the homotopic class γ of the loop $\gamma(x_0, x_0)$. We denote by M_γ the value $Y(x_0^*(\gamma); x_0, I)$ of the solution of the equation (5.1), normalized by the initial condition (5.4), at the point $x_0^*(\gamma)$. The correspondence $\gamma \mapsto M_\gamma$ is a representation⁵ of the fundamental group $\pi(\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}; x_0)$ into the general linear group $GL(k; \mathbb{C})$:

$$M_{\gamma'\gamma''} = M_{\gamma'}M_{\gamma''} \quad \text{for every } \gamma', \gamma'' \in \pi(\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}; x_0)$$

⁵As usual, the product $\gamma'\gamma''$ is defined by the composed loop, obtained by traversing first the loop, generating γ'' , and then the loop, generating γ' .

and

$M_\varepsilon = I$, where ε is the unity of the group $\pi(\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}; x_0)$

Definition 5.2. The matrix M_γ is said to be the monodromy matrix for the equation (5.1), corresponding to the homotopic class γ . The correspondence $\gamma \mapsto M_\gamma$ is said to be the monodromy representation of the equation (5.1).

The group $\pi(\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}; x_0)$ is finitely generated. Let us choose a special system of generators of this group. Every simple loop (i.e. the loop which is homeomorphic to a circle) divides the Riemann sphere $\overline{\mathbb{C}}$ into two components. Let γ_j be a simple loop in the space $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, with the distinguished point x_0 , such that the point t_j and the set of all other points $\{t_p\}_{p \neq j}$ belong to the different components and, moreover, the loop γ_j goes around the point t_j in the counterclockwise direction. (See Figure 1). We denote the homotopic class of this loop by γ_j as well. It is known that the elements $\gamma_1, \dots, \gamma_n$ generate the fundamental group $\pi(\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}; x_0)$ and that the equality $\gamma_1 \cdots \gamma_n = \varepsilon$ is the only generating relation⁶.

For what follows, it is crucial to know how a solution of a Fuchsian differential equation behaves in a neighborhood of its singular point.

More generally, let us discuss the behavior of a solution of the differential equation of the form

$$(5.5) \quad \frac{dY}{dx} = \left(\frac{Q}{x-t} + \Phi(x) \right) Y(x),$$

where Q is a constant $k \times k$ matrix, and Φ is a $k \times k$ matrix function, which is holomorphic in the disk $D_{t,\rho}$, centered at the point $x = t$, of radius ρ . Solutions of the equation (5.5) are holomorphic (in general, multi-valued) functions in the punctured disk $D_{t,\rho} \setminus \{t\}$.

The equation (5.5) can be considered as a perturbation of the equation

$$(5.6) \quad \frac{dY}{dx} = \frac{Q}{x-t} Y(x).$$




FIGURE 1.

⁶The generating relation in this form holds for the loops, arranged as shown in Figure 1. In the general case, this relation is of the form $\gamma_{p_1} \cdots \gamma_{p_n} = \varepsilon$ where $p_1, \dots, p_n$ is a permutation of the indices $1, \dots, n$ which is determined by the arrangement of the loops.

The "unperturbed" equation (5.6) can be solved explicitly. It has a fundamental solution of the form

$$(5.7) \quad Y(x) = (x - t)^Q.$$

(By definition, $(x - t)^Q \stackrel{\text{def}}{=} e^{Q \ln(x-t)}$). In general, the function $(x - t)^Q$ considered as a function in the complex plane is multivalued. It can be considered as a univalued holomorphic matrix function on the universal covering $\text{cov}(\mathbb{C} \setminus \{t\})$ of the complex plane punctured at the point $\{t\}$ (or, in other terminology, on the Riemann surface of $\log(x - t)$). The differentiation rule

$$(5.8) \quad \frac{d}{dx}(x - t)^Q = \frac{Q}{x - t}(x - t)^Q, \quad x \in \text{cov}(\mathbb{C} \setminus \{t\}),$$

holds. (The values of $(x - t)^Q$ in both sides of (5.8) are the same).

The structure of the solution of the original "perturbed" equation (5.5) depends on the nature of the spectrum of the matrix Q .

Definition 5.3. A square matrix Q is said to be **non-resonant** if no two eigenvalues of Q differ by a nonzero integer, or in other words, if the spectra of the matrices $Q + nI$ and Q are disjoint for every $n \in \mathbb{Z} \setminus 0$.

Remark 5.4. It may happen that some eigenvalues of a non-resonant matrix Q coincide (i.e., the eigenvalues of a non-resonant matrix Q may be multiple).

Proposition 5.5. *If Q is a non-resonant (constant with respect to x) matrix, and $\Phi(x)$ is a matrix-function holomorphic in the disk $D_{t,\rho}$, then the differential equation (5.5) has a fundamental solution $Y(x)$ of the form*

$$(5.9) \quad Y(x) = H(x)(x - t)^Q, \quad x \in \text{cov}(D_{t,\rho} \setminus \{t\}),$$

defined on the universal covering $\text{cov}(D_{t,\rho} \setminus \{t\})$ of the punctured disc $D_{t,\rho} \setminus \{t\}$, where $H(x)$ is a matrix function, holomorphic and invertible⁷ in the disk $D_{t,\rho}$.

The proof of Proposition 5.5, and even of a more general statement, is given in Appendix.

In the general case, when the eigenvalues of the matrix Q can differ by non-zero integers, the situation is more complicated.

Proposition 5.6. *The differential equation (5.5), with a constant matrix Q and a matrix function $\Phi(x)$, holomorphic in the disk $D_{t,\rho}$, has a fundamental solution $Y(x)$ of the form*

$$(5.10) \quad Y(x) = H(x)(x - t)^L(x - t)^{\hat{Q}}, \quad x \in \text{cov}(D_{t,\rho} \setminus \{t\}),$$

where $H(x)$ is a matrix function holomorphic and invertible in the disk $D_{t,\rho}$, L is a diagonalizable matrix with integer eigenvalues $l_1, \dots, l_k$, and $\hat{Q}$ is a

⁷The inverse matrix function $H^{-1}(x)$ exists and is holomorphic in $D_{t,\rho}$

non-resonant matrix whose eigenvalues $\hat{\lambda}_1, \dots, \hat{\lambda}_k$ are related to the eigenvalues $\lambda_1, \dots, \lambda_k$ of the matrix Q according to

$$(5.11) \quad \hat{\lambda}_p = \lambda_p - l_p, \quad 1 \leq p \leq k.$$

For a certain choice of the integers $l_1 \geq \dots \geq l_n$, the matrices L and Q can be simultaneously reduced to the forms

$$L = T^{-1} \text{Diag}(l_1, \dots, l_n) T, \quad \hat{Q} = T^{-1} \left(\text{Diag}(\hat{\lambda}_1, \dots, \hat{\lambda}_n) + U \right) T,$$

where U is an upper-triangular matrix with zero diagonal.

Proposition 5.6 is applicable to equation (5.5) with the general matrix Q . In particular, it is applicable to the equation with the non-resonant matrix Q . However, if Q is non-resonant, one can choose $L = 0$ and $\hat{Q} = Q$.

Remark 5.7. It should be mentioned that the matrix function $H(x)$ in (5.12) (or in (5.15)) is not quite unique. The matrix $H(t) = H(x)_{x=t}$ can be an arbitrary invertible matrix commuting with the matrix Q . Of course, the solution (5.9) (or (5.10)) depends on this choice. In the *non-resonant case*, as soon as this choice is made, the local solution of the equation (5.5) at the point $x = t$, i.e. the solution of the form (5.9), is determined uniquely (up to the choice of $\arg(x - t)$ which leads to the non-uniqueness of $(x - t)^Q = e^{Q \ln(x-t)}$). In what follows, we choose

$$H(x)_{|x=t} = I.$$

In the non-resonant case, the latter equality can be considered as a *normalizing condition* for the local solution (5.9) of (5.5) at the point $x = t$.

However, in the resonant case, the condition $H(x)_{|x=t} = I$ do *not* determine the solution of the form (5.10) of the equation (5.5) uniquely. The following example illustrates this non-uniqueness. Consider the differential equation

$$\frac{\partial Y(x)}{\partial x} = \frac{Q}{x} \cdot Y(x),$$

where

$$Q = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$$

This equation, which is of the equation (5.5) with $t = 0, \Phi(x) \equiv 0$, has the solution

$$Y_1(x) = \begin{bmatrix} x & 0 \\ 0 & 1 \end{bmatrix},$$

which is of the form (5.10) with $H(x) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, L = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \hat{Q} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, t = 0$. The matrix function

$$Y_2(x) = Y_1(x) \cdot \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \quad \text{or, more explicitly,} \quad Y_2(x) = \begin{bmatrix} x & x \\ 0 & 1 \end{bmatrix},$$

is another solution of the same differential equation. However, the function $Y_2(x)$ is also representable in the form

$$Y_2(x) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x & 0 \\ 0 & 1 \end{bmatrix},$$

i.e. in the form (5.10), with $H(x) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$, $L = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, $\hat{Q} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, $t = 0$.

Thus, the considered differential equation possesses two different solutions $Y_1(x)$ and $Y_2(x)$, both these solutions are of the form (5.10), with the same L and $\hat{Q}$, but with different $H(x)$, and both these factors $H(x)$ satisfy the condition $H(x)|_{x=0} = I$.

Thus, *in the resonant case, the condition $H(x)|_{x=t} = I$ does not uniquely determine the local solution (5.10) of the equation (5.5) corresponding to the singular point $x = t$, even though the matrices L and $\hat{Q}$ are specified.*

Remark 5.8. There is another essential difference between the non-resonant and the "resonant" cases. If the matrix Q is not non-resonant, then the matrix $\hat{Q}$, which appears in (5.10), depends on the value $\Phi(t) = \Phi(x)|_{x=t}$ of the "perturbing" (see (5.5)) function Φ at the point t . In particular, the Jordan structure of the matrix $\hat{Q}$ may depend on the matrix $\Phi(t)$. This Jordan structure, together with the eigenvalues of $\hat{Q}$, determine the multi-valued behavior of the matrix function $(x - t)^{\hat{Q}}$, and finally, the multi-valued behavior of $Y(x)$ in $D_{t,\rho}$. Thus, in the "resonant" case the multi-valued behavior of the solution $Y(x)$ depends not only on the matrix Q (the coefficient of the "leading term") but also on the "perturbing" matrix Φ . As can be seen from (5.9), in the non-resonant case the multi-valued behavior of the solution $Y(x)$ depends only on Q , but not on Φ .

Remark 5.9. Of course, for every matrices A and B , the matrix $e^A \cdot e^B$ can be presented as e^C :

$$e^A \cdot e^B = e^C, \quad \text{where} \quad C = \log(e^A \cdot e^B),$$

where an arbitrary value for the matrix $\log(e^A \cdot e^B)$ can be chosen. However, the case of exponential functions is different. In general, a matrix function $e^{A\zeta} \cdot e^{B\zeta}$ of ζ (with constant matrices A and B) is not representable as $e^{C\zeta}$ with a constant matrix C . In particular, in the resonant case the solution (5.10) of the differential equation (5.5) is not, in general, representable in the form $Y(x) = H(x) \cdot (x - t)^R$ with a holomorphic invertible matrix function H and a constant matrix R .

Propositions 5.5 and 5.6 go back to L. Fuchs and G. Frobenius. The systematic study of linear differential equations with analytic coefficients in the neighborhood of a singular point was initiated by Lazarus Fuchs in 1866 and followed by G. Frobenius in 1873. They discussed the "scalar" differential equation of the n^{th} order. Matrix differential equations in the complex domain were considered later. The statement which we formulated

as Proposition 5.5 can be found in a lot of excellent textbooks now. See, for example, [CoLe], Theorem 4.1; [Gant], Chapter XV, §10; [Hart], Chapter IV, Corollary 11.2; [IKSY], Chapter I, Theorem 2.3. However, it is difficult to point out the source, where this statement first appeared precisely in the same form as in Proposition 5.5. The earliest reference we know is [Ra] (see Section 3 there). The statement which we formulated as Proposition 5.6 can be found in [Gant], Chapter XV, §10, in [HsSi], Chapter V, Section 5; [Sib], Chapter III, Theorem 3.9.4, page 89. The earliest of these references is [Gant] (the first edition of Gantmacher's book appeared in 1954). However, Y. Sibuya, [Sib], page 89, refers to Hukuhara's book [Huk] (Theorem 41.1, page 143) published in 1950. Unfortunately, the book [Huk] is unavailable to us.

Let us return to the Fuchsian equation (5.1). We consider this equation in a disk neighborhood $D_{t_j, \rho}$ of the singular point t_j , which does not contain other singular points $t_p, p \neq j$. Let us choose and fix a point x_j of the punctured disk $D_{t_j, \rho} \setminus \{t_j\}$. We also choose and fix a path α_j in $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, which starts at the distinguished point x_0 and ends at the




FIGURE
2. a:
path α_j .




FIGURE
2. b:
path μ .




FIGURE
2. c:
path α .

point x_j . (See Figure 2.a.) We need these x_j and α_j to distinguish a certain analytic continuation Y_{α_j} of the solution of (5.1) normalized by (5.4) into the punctured disk $D_{t_j, \rho} \setminus \{t_j\}$. Let x be an arbitrary point of the punctured disk $D_{t_j, \rho} \setminus \{t_j\}$, and μ be an arbitrary path in $D_{t_j, \rho} \setminus \{t_j\}$, starting at x_j and ending at x . (See Figure 2.b.). So, the pair (x, μ) can be considered as a point of the universal covering $\text{cov}(D_{t_j, \rho} \setminus \{t_j\}; x_j)$. We construct the composed path $\alpha = \mu \cdot \alpha_j$. The path α starts at the point x_0 and ends at the point x . (See Figure 2.c.) We continue analytically the solution Y along the path α from the point x_0 to the point x . In this manner we distinguish

the continuation $Y_{\alpha_j}(x)$ of the solution $Y(\cdot)$, normalized by the condition (5.4), from a neighborhood of x_0 to the punctured disk $D_{t_j, \rho} \setminus \{t_j\}$. The function $Y_{\alpha_j}(\cdot)$ can be considered as a multi-valued function in $D_{t_j, \rho} \setminus \{t_j\}$, or, better to say, as a single-valued function in the universal covering $\text{cov}(D_{t_j, \rho} \setminus \{t_j\}; x_j)$ with the distinguished point x_j . The notation Y_{α_j} reflects how the path α_j is involved in the process of the analytic continuation. In fact, this process involves two steps. On the first step, the solution is continued along the path α_j from a neighborhood of the initial point x_0 to a neighborhood of the point x_j , where the point x_j is considered as a distinguished point of the punctured disk $D_{t_j, \rho} \setminus \{t_j\}$. On the second step, the solution is continued along paths which are contained in $D_{t_j, \rho} \setminus \{t_j\}$ from the neighborhood of the point x_j to points of $\text{cov}(D_{t_j, \rho} \setminus \{t_j\}; x_j)$.

The matrix function $Y_{\alpha_j}(\cdot)$ is a fundamental solution of (5.1). Propositions 5.5-5.6 provide another fundamental solution of (5.1). As a preliminary we assume that the matrix Q_j is non-resonant. According to Proposition 5.5, there exists the fundamental solution $Y_j(x)$ of (5.1), which is representable as

$$(5.12) \quad Y_j(x) = H_j(x)(x - t_j)^{Q_j}, \quad x \in \text{cov}(D_{t_j, \rho} \setminus \{t_j\}, x_j),$$

where $H_j(x)$ is a matrix function, holomorphic and invertible in the whole (non-punctured) disc $D_{t_j, \rho}$. Under the normalizing condition

$$(5.13) \quad H_j(x)|_{x=t_j} = I,$$

the matrix function $H_j(x)$ is unique. The matrix function $(x - t_j)^{Q_j}$ is defined up to a right factor of the form $\exp[2\pi i Q_j]$. To avoid this non-uniqueness, we choose a certain value

$$(5.14) \quad \theta_j = \arg(x_j - t_j),$$

and fix this choice. Under the agreement (5.14), the matrix function $(x - t)^{Q_j}$ is defined uniquely on $\text{cov}(D_{t_j, \rho} \setminus \{t_j\}, x_j)$. Thus, *if the matrix Q_j is non-resonant, the local solution $Y_j(x)$ of the differential equation (5.1) corresponding to the singular point t_j can be distinguished uniquely by means of the normalizing conditions (5.13) and (5.14).*

If the matrix Q_j is resonant, the situation is different. According to Proposition 5.6, there exists the fundamental solution $Y_j(x)$ of (5.1), which is representable as

$$(5.15) \quad Y_j(x) = H_j(x)(x - t_j)^{L_j}(x - t_j)^{\hat{Q}_j}, \quad x \in \text{cov}(D_{t_j, \rho} \setminus \{t_j\}),$$

where the matrix $H_j(x)$ has the same properties as before, L_j is a diagonalizable matrix with integer eigenvalues (so, the matrix function $(x - t_j)^{L_j}$ is single-valued in $D_{t_j, \rho} \setminus \{t_j\}$), and $\hat{Q}_j$ is a non-resonant matrix, whose eigenvalues differ from eigenvalues of the matrix Q_j by some integer numbers. However, even though the normalizing conditions (5.13) and (5.14) are assumed and the matrices $L_j, \hat{Q}_j$ are chosen, the matrix function $H_j(x)$ in

(5.15)) is determined *non-uniquely*⁸. Thus, if the matrix Q_j is resonant, we may deal with some local solution $Y_j(x)$ of the form (5.15) corresponding to the singular point t_j of the differential equation (5.1), but in general, we can not distinguish naturally such a solution.

Definition 5.10. Assume that the matrix Q_j is non-resonant. The solution Y_j of the Fuchsian equation (5.1) representable in the form (5.12) near $\{t_j\}$, with H_j satisfying the normalizing condition (5.13), is said to be the normalized local solution of (5.1) corresponding to the singular point $\{t_j\}$.

Let us choose some local solution Y_j of the equation (5.1) corresponding to the singular point t_j . If the matrix Q_j is non-resonant, it is natural to choose the normalized local solution. If the matrix Q_j is not non-resonant, we choose some local solution of the form (5.10) and fix such a choice. Since both solutions, Y_{x_j, α_j} and Y_j , are fundamental, the equality

$$(5.16) \quad Y_{x_j, \alpha_j}(x) = Y_j(x)C,$$

holds, where C is a constant (with respect to x) non-degenerate matrix.

Definition 5.11. The equality (5.16) is said to be the connection relation. The matrix C from (5.16) is said to be the connection coefficient.

According to Definition 5.11, the connection coefficient C depends on the choice of the point $x_j \in D_{t_j, \rho} \setminus \{t_j\}$ and on the choice of the path α_j , connecting the points x_0 and x_j : $C = C_{x_j, \alpha_j}$. Now we will discuss this dependence.

Let α'_j and α''_j be two paths, both of which start at the point x_0 and end at the point x_j , $\alpha'_j, \alpha''_j \subset \overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$. It is clear that $\alpha''_j = \alpha'_j \cdot \gamma$, where γ is the loop with the distinguished point x_0 , which is composed of the path α''_j and of the path $(\alpha'_j)^{-1}$ (the latter is the path α'_j which is traversed in the opposite direction: from the point x_j to the point x_0): $\gamma = (\alpha'_j)^{-1} \cdot \alpha''_j$. The paths α' and α'' , connecting the point x_0 with the point x , are defined as the compositions $\alpha' = \mu \cdot \alpha'_j$ and $\alpha'' = \mu \cdot \alpha''_j$, where, as before, μ , $\mu \subset D_{t_j, \rho} \setminus \{t_j\}$, is a path connecting the points x_j and x . Thus, $\alpha'' = \alpha' \cdot \gamma$, and

$$Y_{\alpha''_j}(x) = Y_{\alpha'_j}(x)M_\gamma,$$

where M_γ is the monodromy matrix corresponding to the loop γ . On the other hand,

$$Y_{\alpha'_j}(x) = Y_j(x)C_{x_j, \alpha'_j}, \quad Y_{\alpha''_j}(x) = Y_j(x)C_{x_j, \alpha''_j}.$$

Therefore,

$$(5.17) \quad C_{x_j, \alpha''_j} = C_{x_j, \alpha'_j}M_\gamma, \quad \text{where } \gamma = (\alpha'_j)^{-1} \cdot \alpha''_j.$$

⁸See Remark 5.7. In what follows, such an arbitrariness in the choice of the local solution does not play any essential role. We need the notions of the local solution and of the connection coefficient only to *illustrate* and *motivate* the notion of the isoprincipal deformation which will be introduced further, in section 7.

Let us discuss, how the connection coefficient C depends on the point x_j . Given two points $x'_j, x''_j \in D_{t_j, \rho} \setminus \{t_j\}$ and a path $\alpha'_j \subset \overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, connecting x_0 and x'_j , there exists a path $\alpha''_j \subset \overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, connecting x_0 and x''_j , such that

$$(5.18) \quad C_{x'_j, \alpha'_j} = C_{x''_j, \alpha''_j},$$

Indeed, let ν be a path, connecting x'_j and x''_j , $\nu \subset D_{t_j, \rho} \setminus \{t_j\}$. We define the path α''_j connecting x_0 with x''_j as the composition $\alpha''_j = \nu \cdot \alpha'_j$. The pairs (x'_j, α'_j) and (x''_j, α''_j) allow us to distinguish the solutions $Y_{\alpha'_j}$ and $Y_{\alpha''_j}$ of the equation (5.1) in the punctured neighborhood $D_{t_j, \rho} \setminus \{t_j\}$. To obtain values of these solutions at a point $x \in D_{t_j, \rho} \setminus \{t_j\}$, we have to choose paths μ'_j and μ''_j , connecting the points x'_j and x''_j , respectively, with the point x . If we make this choice coherently, we obtain the equality $Y_{\alpha'_j}(x) = Y_{\alpha''_j}(x)$ which implies (5.18). Such a coherent choice can be done in the following way. If μ' is a path in $D_{t_j, \rho} \setminus \{t_j\}$, connecting x'_j with x , we define the path μ'' , connecting x''_j with x , as the composition $\mu'' = \mu' \cdot \nu^{-1}$. To obtain the values of the solutions $Y_{x'_j, \alpha'_j}$ and $Y_{x''_j, \alpha''_j}$ at the point x , we have to continue analytically the solution Y , which is normalized by (5.4), from the point x_0 to the point x along the paths $\alpha' = \mu' \cdot \alpha'_j$ and $\alpha'' = \mu'' \cdot \alpha''_j$, respectively. However, these two paths are homotopic. To see this, we remark that $\alpha'' = (\mu' \cdot \nu^{-1}) \cdot (\nu \cdot \alpha') = \mu' \cdot (\nu^{-1} \cdot \nu) \cdot \alpha'$ and that the path $\nu^{-1} \cdot \nu$ is contractible to the point x'_j . Hence, $Y_{\alpha'_j}(x) = Y_{\alpha''_j}(x)$. Thus, in some sense the connection coefficient does not depend on the choice of the point $x_j \in D_{t_j, \rho} \setminus \{t_j\}$.

Let us calculate the monodromy matrix corresponding to the loop “going around” the point t_j . We relate such a loop γ_{α_j} to every path α_j , $\alpha_j \subset \overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, which starts at the point x_0 and ends at a point x_j belonging to a punctured disk $D_{t_j, \rho} \setminus \{t_j\}$. The radius ρ must be chosen so small that the disk does not contain any other point $t_p, p \neq j$. Let μ be the circular arc, centered at the point t_j , of the radius $\varepsilon = |x_j - t_j|$, which starts at the point x_j , ends at the point $t_j + (x_j - t_j)e^{2\pi i}$ and is oriented counterclockwise, making precisely one turn around its center t_j . We construct the loop γ_{α_j} as the composition of the path α_j , of the arc μ and of the path α_j^{-1} : $\gamma = (\alpha_j)^{-1} \cdot \mu \cdot \alpha_j$ (see Figure 3).




FIGURE 3. Loop γ_{α_j} .

Definition 5.12. The loop γ_{α_j} is said to be the loop generated by the path α_j .

Lemma 5.13. *The monodromy matrix $M_{\gamma_{\alpha_j}}$ along the loop γ_{α_j} generated by a path α_j can be represented as*

$$(5.19) \quad M_{\gamma_{\alpha_j}} = (Y_{\alpha_j}(x_j))^{-1} Y_{\alpha_j}(t_j + (x_j - t_j)e^{2\pi i}),$$

where x_j is the endpoint of the path α_j .

PROOF. For linear differential equations, the following *semigroup property* holds: Let γ be a path starting at a point a and ending at the point c , and let b is an “intermediate” point of the path α , that is the point which is located “between” the points a and c . Then $Y(c, a) = Y(c, b)Y(b, a)$, where for $\xi, \eta \in \alpha$, $Y(\eta, \xi)$ is the value $Y(x, \xi)|_{x=\eta}$ of the solution of the equation satisfying the initial condition $Y(x, \xi)|_{x=\xi} = I$. Let us take the loop γ_{α_j} as such a path γ . The point a is the point x_0 considered as the starting point of the loop, the point c is the same point x_0 , but considered as the end point of the loop. The point b is the point $t_j + (x_j - t_j)e^{2\pi i}$. From the semigroup property it follows that $Y(b, c)Y(c, b) = I$. It is clear that $Y(b, a) = Y_{\alpha_j}(t_j + (x_j - t_j)e^{2\pi i})$, $Y(b, c) = Y_{\alpha_j}(x_j)$, and $Y(c, a) = M_{\gamma_{\alpha_j}}$. Q.E.D.

From (5.16) it follows that in the non-resonant case,

$$Y_{x_j, \alpha_j}(x_j) = H_j(x_j)(x_j - t_j)^{Q_j} C_{\alpha_j}$$

and

$$Y_{x_j, \alpha_j}(t_j + (x_j - t_j)e^{2\pi i}) = H_j(x_j)e^{2\pi i Q_j}(x_j - t_j)^{Q_j} C_{\alpha_j}.$$

(The function $H_j(x)$ is single-valued in $D_{t_j, \rho}$, and the function $(x - t_j)^{Q_j}$ receives the factor $e^{2\pi i Q_j}$, when x traverses the arc μ once, in the counter-clockwise direction.) Therefore, according to Lemma 5.13,

$$(5.20) \quad M_{\gamma_j} = C_{\alpha_j}^{-1}(x_j - t_j)^{-Q_j} e^{2\pi i Q_j}(x_j - t_j)^{Q_j} C_{\alpha_j}.$$

Finally, we obtain

Lemma 5.14. *The monodromy matrix $M_{\gamma_{\alpha_j}}$, corresponding to the loop γ_{α_j} generated by a path α_j can be represented as*

$$(5.21) \quad M_{\gamma_j} = C_{\alpha_j}^{-1} \exp[2\pi i Q_j] C_{\alpha_j}$$

in the non-resonant case, and as

$$(5.22) \quad M_{\gamma_j} = C_{\alpha_j}^{-1} \exp[2\pi i \hat{Q}_j] C_{\alpha_j}$$

in the resonant case, where C_{α_j} is the connection coefficient, corresponding to the path α_j .

(The reasoning in the resonant case is analogous. One needs to use the equality (5.15), taking into account that the matrix-function $(x - t_j)^{L_j}$ is single-valued in the disk $D_{t_j, \rho}$).

For what follows, it is very fruitful to rewrite the representations (5.12) - (5.16), (5.15) - (5.16) for a “local” solution ⁹of (5.1), as well as the expressions (5.21) and (5.22) for the monodromy matrices, in a different form. The respective transformations are trivial, but nevertheless very useful. The form in which we present the local solutions hints at how to introduce the notion of the isoprincipal deformation of the Fuchsian differential equation. This notion will be of crucial importance to us.

Let $D_{t_j, \rho}$ be a disk neighborhood of the singular point t_j of the differential equation (5.1), which is small enough, so that the point t_j is the only singular point which is contained in this neighborhood. In this neighborhood lives a “local” solution Y_j of the equation (5.1), which in the non-resonant case is of the form (5.12), and in the resonant case is of the form (5.15). However, this “local” solution Y_j does not, in general, satisfy the initial condition (5.4). To be precise, we should speak about the analytic continuation of this solution from the neighborhood $D_{t_j, \rho}$ to the point x_0 . To discuss such an analytic continuation, we have to choose a point x_j of the punctured neighborhood $D_{t_j, \rho} \setminus \{t_j\}$, and a path α_j , $\alpha_j \subset \overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, leading from the point x_0 to the point x_j . The solution Y_{α_j} which satisfies the initial condition (5.4) at the distinguished point x_0 and which is continued analytically to the neighborhood $D_{t_j, \rho}$ along the path α_j is of the form

$$(5.23) \quad Y_{\alpha_j}(x) = H_j(x)(x - t_j)^{Q_j} C_{\alpha_j}, \quad x \in \text{cov}(D_{t_j, \rho} \setminus \{t_j\}, x_j),$$

in the non-resonant case, and of the form

$$(5.24) \quad Y_{\alpha_j}(x) = H_j(x)(x - t_j)^{L_j}(x - t_j)^{\hat{Q}_j} C_{\alpha_j}, \quad x \in \text{cov}(D_{t_j, \rho} \setminus \{t_j\}, x_j),$$

in the resonant case, where C_{α_j} is the connection coefficient corresponding to the choice of the homotopic class of the path α_j and the choice of the arguments $\arg(x_j - t_j)$, see (5.14), (5.12) and (5.15).

We introduce the matrices

$$(5.25) \quad A_{\alpha_j} = C_{\alpha_j}^{-1} Q_j C_{\alpha_j}, \quad H_{\alpha_j} = H_j C_{\alpha_j}$$

in the non-resonant case, and the matrices

$$(5.26) \quad A_{\alpha_j} = C_{\alpha_j}^{-1} \hat{Q}_j C_{\alpha_j}, \quad B_{\alpha_j} = C_{\alpha_j}^{-1} L_j C_{\alpha_j}, \quad H_{\alpha_j} = H_j C_{\alpha_j}$$

in the resonant case. Propositions 5.5 and 5.6 can be “developed” in the following way.

Theorem 5.15. *If the matrix Q_j is non-resonant, then the solution Y_{α_j} of the differential equation (5.1) that satisfies the initial condition (5.4) at the distinguished point x_0 and is continued analytically along a path α_j into a small punctured neighborhood $D_{t_j, \rho} \setminus \{t_j\}$ of the singular point t_j , is representable in the form*

$$(5.27) \quad Y_{\alpha_j}(x) = H_{\alpha_j}(x)(x - t_j)^{A_{\alpha_j}}, \quad x \in \text{cov}(D_{t_j, \rho} \setminus \{t_j\}, x_j),$$

⁹I.e., for a solution in a neighborhood of the singular point t_j .

where A_{α_j} is a matrix, similar to the matrix Q_j , and H_{α_j} is a matrix function, holomorphic and invertible in the entire neighborhood $D_{t_j,\rho}$ (including the point t_j).

Theorem 5.16. *If the matrix Q_j is resonant, then the solution Y_{α_j} of the differential equation (5.1), that satisfies the initial condition (5.4) at the distinguished point x_0 and is continued analytically along a path α_j into a small punctured neighborhood $D_{t_j,\rho} \setminus \{t_j\}$ of the singular point t_j , is representable in the form*

$$(5.28) \quad Y_{\alpha_j}(x) = H_{\alpha_j}(x)(x-t_j)^{Z_{\alpha_j}}(x-t_j)^{A_{\alpha_j}}, \quad x \in \text{cov}(D_{t_j,\rho} \setminus \{t_j\}, x_j),$$

where Z_{α_j} is a diagonalizable matrix with integer eigenvalues $l_1, \dots, l_k$, A_{α_j} is a non-resonant matrix, whose eigenvalues $\hat{\lambda}_p$ are related to the eigenvalues λ_p of the matrix Q_j by the relation

$$(5.29) \quad \hat{\lambda}_p = \lambda_p - l_j, \quad 1 \leq p \leq k,$$

and $H_{\alpha_j}(x)$ is a matrix function, holomorphic and invertible in the entire neighborhood (including the point t_j). For a certain choice of the integers $l_1 \geq \dots \geq l_n$, the matrices Z_{α_j} and A_{α_j} can be simultaneously reduced to the forms

$$Z_{\alpha_j} = T_{\alpha_j}^{-1} \text{Diag}(l_1, \dots, l_n) T_{\alpha_j}, \quad A_{\alpha_j} = T_{\alpha_j}^{-1} (\text{Diag}(\hat{\lambda}_1, \dots, \hat{\lambda}_n) + U) T_{\alpha_j},$$

where U is an upper-triangular matrix with zero diagonal.

Remark 5.17. If α'_j is another path with the same starting point x_0 and endpoint x_j , then, in the non-resonant case, the matrices A_{α_j} and H_{α_j} from (5.27) are transformed according to the rule

$$(5.30) \quad A_{\alpha'_j} = M_{\gamma}^{-1} A_{\alpha_j} M_{\gamma}, \quad H_{\alpha'_j}(x) = H_{\alpha_j}(x) M_{\gamma},$$

and in the resonant case, the matrices Z_{α_j} , A_{α_j} and H_{α_j} from (5.28) are transformed according to the rule

$$(5.31) \quad Z_{\alpha'_j} = M_{\gamma}^{-1} Z_{\alpha_j} M_{\gamma}, \quad A_{\alpha'_j} = M_{\gamma}^{-1} A_{\alpha_j} M_{\gamma}, \quad H_{\alpha'_j}(x) = H_{\alpha_j}(x) M_{\gamma},$$

where M_{γ} is the monodromy matrix corresponding to the loop $\gamma = (\alpha_j)^{-1} \alpha'_j$. In particular, if the paths α_j and α'_j are homotopic then

$$Z_{\alpha'_j} = Z_{\alpha_j}, \quad A_{\alpha'_j} = A_{\alpha_j}, \quad H_{\alpha'_j}(x) = H_{\alpha_j}(x).$$

Theorem 5.18. *Let $M_{\gamma_{\alpha_j}}$ be the monodromy matrix for the differential equation (5.1) corresponding to the loop γ_{α_j} , which goes around the singular point t_j and is constructed from the path α_j , connecting the initial point x_0 to a small neighborhood of t_j : $\gamma_j = (\alpha_j)^{-1} \cdot \mu \cdot \alpha_j$ (see Figure 3). Then this monodromy matrix can be expressed as*

$$(5.32) \quad M_{\gamma_{\alpha_j}} = \exp(2\pi i A_{\alpha_j}),$$

where the matrix A_{α_j} is the same as in the exponent in (5.27), if Q_j is non-resonant, and the exponent in (5.28), if Q_j is resonant.

Theorem 5.19. *Let M_{γ_j} be the monodromy matrix for the differential equation (5.1) corresponding to a loop γ_j with the distinguished point x_0 which makes one turn counterclockwise around the singular point t_j and makes no turns around the other singular points $t_p, p = 1, \dots, t_n, p \neq j$. Then, whatever the matrix Q_j is, non-resonant or not, the spectra $\text{spec } M_{\gamma_j}$ and $\text{spec } Q_j$ are related by*

$$(5.33) \quad \text{spec } M_{\gamma_j} = \exp \{2\pi i \text{spec } Q_j\},$$

where the latter equality holds “with multiplicities”, that is the multiplicity κ_μ of the eigenvalue μ of the matrix M_{γ_j} and multiplicities κ_λ of the eigenvalues λ of the matrix Q_j are related by $\kappa_\mu = \sum \kappa_\lambda$, where the sum is taken over all λ such that $\exp\{\lambda\} = \mu$.

PROOF. This theorem is the immediate consequence of Theorem 5.18, of the spectral mapping theorem for matrices (from (5.32) it follows that $\text{spec } M_{\gamma_j} = \exp \{2\pi i \text{spec } A_{\alpha_j}\}$), and of Theorems 5.15 or 5.16, which relate the spectra of the matrices A_{α_j} and Q_j . However, here we can do without Theorem 5.16, restricting ourselves to non-resonant matrices Q_j only. Every matrix can be approximated by a non-resonant one. The spectrum $\text{spec } Q_j$ depends on Q_j continuously. The monodromy matrix M_{γ_j} depends on Q_j continuously as well. Q.E.D.

6. DEFORMATIONS OF FUCHSIAN DIFFERENTIAL EQUATIONS

A *deformation* of a Fuchsian equation is a *family* of Fuchsian differential equations depending on parameters:

$$\frac{dY}{dx} = \left(\sum_{1 \leq j \leq n} \frac{Q_p(\alpha)}{x - t_p} \right) Y.$$

If the coefficients $Q_p(\alpha)$ depend on the parameters $\alpha = (\alpha_1, \dots, \alpha_m)$ in some open set of the space $\mathbb{C}^m$ holomorphically, such a deformation is said to be a *holomorphic deformation of a Fuchsian differential equation*. The study of linear differential equations that depend on parameters holomorphically was started by Lazarus Fuchs, see [FuL1] - [FuL4]. This study was continued by L. Schlesinger, see [Sch1] - [Sch3], and by Richard Fuchs, see [FuR]. In particular, L. Schlesinger has considered linear equation, where loci $t_1, \dots, t_n$ of singular points serve as the parameters α . In other words, L. Schlesinger has considered deformations of Fuchsian differential equations of the form

$$(6.1) \quad \frac{dY}{dx} = \left(\sum_{1 \leq p \leq n} \frac{Q_p(\mathbf{t})}{x - t_p} \right) Y,$$

where the coefficients $Q_p(t_1, \dots, t_n), p = 1, \dots, n$, are $k \times k$ matrix functions which are defined and holomorphic for $\mathbf{t}$ from a domain $\mathcal{D}$ of the space

$\mathbb{C}_*^n$. In the further we assume that the condition

$$\sum_{1 \leq p \leq n} Q_p(\mathbf{t}) \equiv 0, \quad \mathbf{t} \in \mathcal{D},$$

is satisfied.

In what follows we study the normalized by (5.4) solution $Y(x, \mathbf{t})$ of the family (6.1) of Fuchsian equations as a function of x and $\mathbf{t}$. In particular, we consider the monodromy matrices M_{γ_j} for the equation (6.1), as well as the matrices A_{α_j} and functions $H_{\alpha_j}(x)$ from the representations (5.27) and (5.28) for various $t_1, \dots, t_n$: $M_{\gamma_j} = M_{\gamma_j}(\mathbf{t})$, $A_{\alpha_j} = A_{\alpha_j}(\mathbf{t})$ and $H_{\alpha_j} = H_{\alpha_j}(\mathbf{t})$, intending to study how these matrices depend on $\mathbf{t} = (t_1, \dots, t_n)$. Here a problem arises. The matrices M_{γ_j} , A_{α_j} , H_{α_j} depend on the homotopy classes of γ_j and α_j , which are considered on the punctured Riemann sphere $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$. However, for different $\mathbf{t} = (t_1, \dots, t_n)$, the sets $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$ are different. Thus we need to explain what does it mean that, for the deformation (6.1) of the Fuchsian equation, the monodromy matrices M_{γ_j} , or the exponents A_{α_j} from (5.27) (or (5.28)) are holomorphic with respect to $\mathbf{t}$, or what does it mean that these matrices do not depend on $\mathbf{t}$. The latter problems are of the local nature.

Let a point $\mathbf{t}^0 = (t_1^0, \dots, t_n^0) \in \mathbb{C}_*^n$ be fixed, $t_j^0 \neq x_0$, $j = 1, \dots, n$, where $x_0 \in \overline{\mathbb{C}}$ is the distinguished point (the same point which appears in the initial condition (5.4), and which serves as the distinguished point for the fundamental group $\pi(\overline{\mathbb{C}} \setminus \{t_1^0, \dots, t_n^0\}; x_0)$). Choose and fix simple loops $\gamma_j, j = 1, \dots, n$, each of which starts and ends at the point x_0 , the loop γ_j contains the point t_j^0 "inside" and goes around this point counterclockwise, and all other points $\{t_p^0\}_{p \neq j}$ lie "outside" the loop γ_j , $j = 1, \dots, n$. Let $\delta > 0$ be a small number such that the disk $D_{t_j^0, \delta} = \{z \in \mathbb{C} : |z - t_j^0| < \delta\}$ does not intersect the loop $\gamma_j, j = 1, \dots, n$ (and hence, is contained inside the loop γ_j). Then for every $\mathbf{t} = (t_1, \dots, t_n)$ such that $t_j \in D_{t_j^0, \delta}$, the loops $\gamma_j, j = 1, \dots, n$, generate the fundamental group $\pi(\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}; x_0)$. See Figure 4, where the disks $D_{t_j^0, \delta}$ are plotted in the light gray color.




FIGURE 4.

Assume that the point $\mathbf{t}^0$ belongs to the domain $\mathcal{D}$ of the space $\mathbb{C}_*^n$, where the matrix functions $Q_j(\mathbf{t})$ are holomorphic, and that δ is so small that the polydisk

$$(6.2) \quad D(\mathbf{t}^0, \delta) = \prod_{p=1}^n D_{t_p^0, \delta}, \quad \text{where} \quad D_{t_p^0, \delta} = \{t_p : |t_p - t_p^0| < \delta\},$$

is contained in $\mathcal{D}$: $D(\mathbf{t}^0, \delta) \subset \mathcal{D}$. Then, for $x \in \gamma_j$, the coefficient matrix $\sum_{1 \leq k \leq n} \frac{Q_j(\mathbf{t})}{x - t_j}$ of the linear equation (6.1) is holomorphic with respect to $\mathbf{t}$ in $D(\mathbf{t}^0, \delta)$. From the standard results on dependence of solution of a linear differential equation on parameters follows

Lemma 6.1. *Under above stated assumptions and notation, the monodromy matrices $M_{\gamma_j}(\mathbf{t})$, $j = 1, \dots, n$, of the holomorphic family (6.1) of Fuchsian equations are holomorphic with respect to $\mathbf{t}$ from the polydisk $D(\mathbf{t}^0, \delta)$.*

Now we investigate in more detail the local solution $Y_j(x, \mathbf{t})$ of the equation (6.1) which corresponds to a neighborhood of the point t_j . We have already considered such a local solution (see Propositions 5.5 - 5.6 and Theorems 5.15 - 5.16). However, until now we considered $\mathbf{t}$ as *fixed* and focused our attention on the multiplicative representation (5.12 - (5.15) of the solution *considered as a function of x* . Now we emphasize the dependence of the local solution on $\mathbf{t}$, while concentrating on the *non-resonant case*. Given $\mathbf{t} \in \mathbb{C}_*^n$ (such that the coefficients $Q_p(\mathbf{t})$, $p = 1, \dots, n$, are defined for this $\mathbf{t}$), and assuming that the matrix $Q_j(\mathbf{t})$ is non-resonant for certain j , the local solution $Y_j(x, \mathbf{t})$ of the equation (6.1) corresponding to the singular point t_j is sought in the form

$$(6.3) \quad Y_j(x, \mathbf{t}) = H_j(x, \mathbf{t})(x - t_j)^{Q_j(\mathbf{t})}, \quad x \in D_{t_j, \rho} \setminus \{t_j\}.$$

The differentiation rule (5.8) for $(x - t_j)^Q$ leads to the differential equation for $H_j(x, \mathbf{t})$:

$$(6.4) \quad \frac{dH_j(x, \mathbf{t})}{dx} = \left(\frac{Q_j(\mathbf{t})H_j(x, \mathbf{t}) - H_j(x, \mathbf{t})Q_j(\mathbf{t})}{x - t_j} + \Phi_j(x, \mathbf{t}) \right) H_j(x, \mathbf{t}),$$

where

$$(6.5) \quad \Phi_j(x, \mathbf{t}) = \sum_{\substack{1 \leq p \leq n, \\ p \neq j}} \frac{Q_p(\mathbf{t})}{x - t_p},$$

the function $\Phi_j(x, \mathbf{t})$ is holomorphic with respect to x at $x = t_j$. The equation (6.4) does not depend on the choice of branch of the (multivalued, in general) function $(x - t_j)^{Q_j(\mathbf{t})}$. The right hand side of this equation, considered as a function of x , has the singularity at the point $x = t_j$. Nevertheless, it can be shown that, if the matrix $Q_j(\mathbf{t})$ is non-resonant, then, under the normalization condition

$$(6.6) \quad H_j(x, \mathbf{t})|_{x=t_j} = I,$$

the holomorphic solution $H_j(x, \mathbf{t})$ of the equation (6.4) exists for x from a disk centered at t_j , and such a solution is unique. More precise formulation of this result will be done below.

In fact, here we repeat what we have already formulated in the Proposition 5.5 and expressed in the formula (5.12). The only difference between the formulae (6.3) and (5.12) is that the notation in (6.3) reflects the dependence on $\mathbf{t}$ explicitly.

Definition 6.2. Let $\mathbf{t} = (t_1, \dots, t_n) \in \mathbb{C}_*^n$ be given, and assume that the matrix $Q_j(\mathbf{t})$ is non-resonant.

The $k \times k$ matrix function $Y_j(x, \mathbf{t})$, possessing the properties

- i. $Y_j(x, \mathbf{t})$ satisfies the differential equation (6.1) with respect to x in a punctured neighborhood $D_{t_j, \rho} \setminus \{t_j\}$ of the point t_j , ($D_{t_j, \rho} = \{z \in \mathbb{C} : |z - t_j| < \rho\}$, ρ is a positive number);
- ii. $Y_j(x, \mathbf{t})$ admits the factorization of the form (6.3), where the factor $H_j(x, \mathbf{t})$ is a matrix function holomorphic with respect to x in the whole (non-punctured) disk $D_{t_j, \rho}$ and satisfying the normalizing condition (6.6),

is said to be the normalized local solution of the differential equation (6.1), corresponding to the singular point t_j .

Proposition 6.3. Let $Q_p(\mathbf{t})$, $p = 1, \dots, n$, be $k \times k$ matrix functions, which are holomorphic with respect to $\mathbf{t} = (t_1, \dots, t_n)$ for $\mathbf{t}$ from an open set $\mathcal{D}$, $\mathcal{D} \in \mathbb{C}_*^n$. Let $\mathbf{t}^0 = (t_1^0, \dots, t_n^0)$ be a point from $\mathcal{D}$, and let

$$(6.7) \quad \rho_q(\mathbf{t}^0) = \frac{1}{2} \min_{\substack{1 \leq p \leq n \\ p \neq q}} |t_q^0 - t_p^0|, \quad 1 \leq q \leq n.$$

Assume that, for certain j , the matrix $Q_j(\mathbf{t}^0)$ is non-resonant. Then there exists $\delta, \delta > 0$, such that for every $\mathbf{t} = (t_1, \dots, t_n)$ from the polydisk ¹⁰ $D(\mathbf{t}^0, \delta)$, the normalized local solution $Y_j(x, \mathbf{t})$ of the equation (6.1), which corresponds to the singular point t_j , exists for x from the punctured disk ¹¹ $D_{t_j, \rho_j(\mathbf{t}^0)} \setminus \{t_j\}$, and the factor $H_j(x, \mathbf{t})$ in (6.3) is a matrix function holomorphic with respect to $x, \mathbf{t}$ for $x \in D_{t_j, \rho_j(\mathbf{t}^0)}$, $\mathbf{t} \in D(\mathbf{t}^0, \delta)$. Moreover, the matrix $H_j(x, \mathbf{t})$ is invertible for these x and $\mathbf{t}$.

The proof of this proposition will be given in Appendix.

Remark 6.4. Proposition 6.3 can be considered as a version of Proposition 5.5, with the emphasis being placed on dependence on $\mathbf{t}$.

Remark 6.5. Proposition 6.3 belongs to the class of statements of the following nature: if coefficients of a differential equations and a normalizing condition depend on some parameters analytically, then the solution of this equation depends on these parameters analytically as well. However, in

¹⁰The polydisk $D(\mathbf{t}^0, \delta)$ was defined in (6.2).

¹¹ $D_{t_j, \rho_j(\mathbf{t}^0)} = \{z \in \mathbb{C} : |z - t_j| < \rho_j(\mathbf{t}^0)\}$.

Proposition 6.3 the normalizing condition (which can be presented in the form $\lim_{x \rightarrow t_j} Y_j(x, \mathbf{t})(x - t_j)^{-Q_j(\mathbf{t})} = I$) is posed at the singular point t_j . Therefore, Proposition 6.3 is not a consequence of standard general results about analytic dependence of a solution of a differential equation on parameters. The proof makes explicit use of the non-resonance of the matrix $Q_j(\mathbf{t}^0)$ and without this assumption the result fails. This shows that the result is rather delicate.

In section 5 the notion of connection coefficients was introduced. The connection coefficients relate the solution of the Fuchsian equation, which is normalized at the distinguished point x_0 , to its local solutions, which are normalized at the singular points of the equation. However, in section 5 we did not care about the dependence of connection coefficients on $\mathbf{t}$. Now we focus our attention on this dependence, concentrating on the non-resonant case.

Given ¹² $\mathbf{t}^0, \mathbf{t}^0 \in \mathcal{D}$, and given $j, 1 \leq j \leq n$, let us choose $\rho_j(\mathbf{t}^0)$ according to (6.7). Then choose $\delta > 0$, satisfying the condition

$$(6.8) \quad 2\delta < \rho_j(\mathbf{t}^0),$$

and choose and fix the point $x_j \in \mathbb{C}$, satisfying the condition

$$(6.9) \quad \delta < |x_j - t_j^0| < \rho_j(\mathbf{t}^0) - \delta.$$




FIGURE 5. a: disk $D_{t_j^0, \rho_j(\mathbf{t}^0)}$.




FIGURE 5. b: disk $D_{t_j, \rho_j(\mathbf{t}^0)}$.

With such a choice

$$(6.10) \quad x_j \in D_{t_j, \rho_j(\mathbf{t}^0)} \quad \text{for every } t_j \in D_{t_j^0, \delta}.$$

(See Figure 5.a, where the disk $D_{t_j^0, \delta}$ is plotted in the light gray color, and the dashed circle is the boundary of the disk $D_{t_j^0, \rho_j(\mathbf{t}^0)}$). Choose a simple path

¹² $\mathcal{D}$ is the open set in $\mathbb{C}_*^n$ where the coefficients $Q_p, 1 \leq p \leq n$, are defined and holomorphic.

α_j , leading from the distinguished point x_0 to the point x_j and separated from the points $t_p^0, p = 1, \dots, t_n$:

$$(6.11) \quad |z - t_p^0| > \delta, \quad \forall z \in \alpha_j, \quad p = 1, \dots, n.$$

Then fix the choice of the path α_j . (In particular, let the path α_j be independent of $\mathbf{t}$.) From (6.11) it follows that for ¹³ $\mathbf{t} = (t_1, \dots, t_n) \in D(\mathbf{t}^0, \delta)$, the condition $|x - t_p| > 0 \quad \forall x \in \alpha_j, p = 1, \dots, n$, holds. Thus, the function $\sum_{1 \leq p \leq n} \frac{Q_p(\mathbf{t})}{x - t_p}$ is holomorphic with respect to $x, \mathbf{t}$ for x from a neighborhood of the path α_j and for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$. Therefore, the solution $Y(x, \mathbf{t})$ of the differential equation (6.1) that satisfies the initial condition

$$(6.12) \quad Y(x, \mathbf{t})_{x=x_0} = I$$

can be continued analytically along the path α_j from a neighborhood $\mathcal{V}_{x_0}$ of the initial point x_0 to a neighborhood $\mathcal{V}_{x_j}$ of the point x_j . ($\mathcal{V}_{x_j}$ does not depend on $\mathbf{t} \in D(\mathbf{t}^0, \delta)$). We denote by $Y_{\alpha_j}(x, \mathbf{t})$ the function in $\mathcal{V}_{x_j}$ which is defined by means of such an analytic continuation. Since the initial condition in the (regular) point x_0 and the path α_j do not depend on $\mathbf{t}$, then for every $x \in \alpha_j$, the solution $Y_{\alpha_j}(x, \mathbf{t})$ is holomorphic with respect to $\mathbf{t}$ for $x \in \mathcal{V}_{x_j}, \mathbf{t} \in D(\mathbf{t}^0, \delta)$. In particular, the matrix $Y_{\alpha_j}(x_j, \mathbf{t})$, as well as the inverse matrix $Y_{\alpha_j}^{-1}(x_j, \mathbf{t})$, is holomorphic with respect to $\mathbf{t}$ for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$. It is clear that the solutions $Y_{\alpha'_j}(x, \mathbf{t})$ and $Y_{\alpha''_j}(x, \mathbf{t})$ corresponding to the analytic continuations into $\mathcal{V}_{x_j}$ along two different paths α'_j and α''_j (with the same endpoints x_0 and x_j), are related by the equality

$$(6.13) \quad Y_{\alpha''_j}(x, \mathbf{t}) = Y_{\alpha'_j}(x, \mathbf{t}) \cdot M_\gamma(\mathbf{t}),$$

where $M_\gamma(\mathbf{t})$ is the monodromy matrix for the equation (6.1) corresponding to the loop $\gamma, \gamma = (\alpha'_j)^{-1} \alpha''_j$, composed from the paths α'_j, α''_j . (The loop γ does not depend on $\mathbf{t}$, but the equation depends. So, the monodromy matrix $M_\gamma(\mathbf{t})$ can depend on $\mathbf{t}$).

Now we turn our attention to the local solution $Y_j(x, \mathbf{t})$, corresponding to the singular point t_j . Assume that the matrix $Q_j(\mathbf{t}^0)$ is non-resonant. Choosing a smaller δ , if necessary, we can apply Proposition 6.3. According to this proposition, the local solution $Y_j(x, \mathbf{t})$ exists in the punctured disk $D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}$. In particular, it is defined at the point x_j , and even in the neighborhood $\mathcal{V}_{x_j}$, if this neighborhood is small enough. The local solution $Y_j(x, \mathbf{t})$ is the product of two factors: $H_j(x, \mathbf{t})$ and $(x - t_j)^{Q_j(\mathbf{t})}$. Under the normalizing condition (6.6), the first factor is determined uniquely. The second factor is not quite unique. In general, the function $(x - t_j)^{Q_j(\mathbf{t})}$ is a multivalued function of x . This function, and hence the solution $Y_j(x, \mathbf{t})$, are determined up to right factor $\exp\{2\pi i Q_j(\mathbf{t})\}$. To avoid this non-uniqueness, we must come to an agreement on the value of $\arg(x - t_j)$ for some fixed

¹³ $D(\mathbf{t}^0, \delta) = D_{t_1^0, \delta} \times \dots \times D_{t_n^0, \delta}$ is a polydisk.

x , for example, for $x = x_j$. However, now $\mathbf{t} = (t_1, \dots, t_n)$ is not a constant but a variable with values in the polydisk $D(\mathbf{t}^0, \delta)$. In particular, the j -th coordinate t_j takes the values in the disk $D_{t_j^0, \rho}$. In order for the function $(x_j - t_j)^{Q_j(\mathbf{t})}$ to be holomorphic with respect to $\mathbf{t}$, we need to choose $\arg(x_j - t_j)$ *coherently* with respect to t_j .

Let us choose somehow and *fix* the value $\arg(x_j - t_j^0)$, say

$$(6.14) \quad \arg(x_j - t_j^0) = \vartheta_j^0.$$

For $t_j \in D_{t_j^0, \delta}$, choose the value of $\arg(x_j - t_j)$ according to the rule:

$$(6.15) \quad \vartheta_0 - \pi/2 < \arg(x_j - t_j) < \vartheta_0 + \pi/2.$$

The choice (6.15) is possible and unique (see Figure 5.a). With such a choice, the matrix $(x_j - t_j)^{Q_j(\mathbf{t})}$ is *determined uniquely* and is holomorphic with respect to $\mathbf{t}$ for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$. Therefore, the matrix $Y_j(x_j, \mathbf{t}) = H_j(x, \mathbf{t})(x_j - t_j)^{Q_j(\mathbf{t})}$ is *determined uniquely* and is holomorphic with respect to $\mathbf{t}$ for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$.

Moreover, the function $Y_j(x, \mathbf{t})$ — the local solution corresponding to the singular point t_j — is determined uniquely on the universal covering $\text{cov}(D_{t_j, \rho_j(\mathbf{t}^0)} \setminus \{t_j\}, x_j)$ with the distinguished point x_j . The conditions (6.14)-(6.15), together with the condition (6.6), can be considered as normalizing condition for the local solution $Y_j(x, \mathbf{t})$: the condition (6.6) normalized the factor $H_j(x, \mathbf{t})$, the condition (6.15) normalized the factor $(x - t_j)^{Q_j(\mathbf{t})}$ coherently with respect to t_j .

Let us define the connection coefficient $C_{\alpha_j}(\mathbf{t})$ as

$$(6.16) \quad C_{\alpha_j}(\mathbf{t}) = Y_j^{-1}(x_j, \mathbf{t}) \cdot Y_{x_j, \alpha_j}(x_j, \mathbf{t})$$

Since both factors in the right hand side of (6.16) are holomorphic with respect to $\mathbf{t} \in D(\mathbf{t}^0, \delta)$ and invertible, *the matrix $C_{\alpha_j}(\mathbf{t})$ is holomorphic with respect to $\mathbf{t} \in D(\mathbf{t}^0, \delta)$ and invertible as well.* According to (6.13), the connection coefficients $C'_{\alpha_j}(\mathbf{t})$ and $C''_{\alpha_j}(\mathbf{t})$, corresponding to two different paths α'_j and α''_j , are related by the equality

$$(6.17) \quad C_{\alpha''_j}(x, \mathbf{t}) = C_{\alpha'_j}(x, \mathbf{t}) \cdot M_\gamma(\mathbf{t}),$$

where $M_\gamma(\mathbf{t})$ is the monodromy matrix for the equation (6.1) corresponding to the loop γ , $\gamma = (\alpha'_j)^{-1} \alpha''_j$.

Since the matrix $C_{\alpha_j}(\mathbf{t})$ does not depend on x , the product $Y_j(x, \mathbf{t}) \cdot C_{\alpha_j}$ is a solution of the differential equation (6.1) for $x \in D_{t_j, \rho_j(\mathbf{t}^0)}$, in particular, for $x \in \mathcal{V}_{x_j}$. The function $Y_{x_j, \alpha_j}(x, \mathbf{t})$ is also a solution of the differential equation (6.1) for $x \in \mathcal{V}_{x_j}$. Both these solutions coincide at the point x_j . Hence they coincide for all $x \in \mathcal{V}_{x_j}$. Therefore, the product $Y_j(x, \mathbf{t}) \cdot C_{\alpha_j}$ can be considered as the *analytic continuation* of the solution $Y_{x_j, \alpha_j}(x, \mathbf{t})$ from $\mathcal{V}_{x_j}$ to the punctured disk $D_{t_j, \rho_j(\mathbf{t}^0)} \setminus \{t_j\}$ (or, better to say, to the universal covering $\text{cov}(D_{t_j, \rho_j(\mathbf{t}^0)} \setminus \{t_j\}; x_j)$ of this punctured disk with the

distinguished point x_j). In its turn, the solution $Y_{x_j, \alpha_j}(x, \mathbf{t})$, which is defined in $\mathcal{V}_{x_j}$, was obtained by the described above analytic continuation of the solution, “living” in $\mathcal{V}_{x_0}$ and satisfying the initial condition (6.12). In other words, the solution $Y_j(x, \mathbf{t}) \cdot C_{\alpha_j}$ can be considered as a solution which is obtained as the result of the “two step” analytic continuation from $\mathcal{V}_{x_0}$ to $\text{cov}(D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}; x_j)$ of the solution, satisfying the initial condition (6.12). However, it is better to talk not about the *two-step* analytic continuation but about the analytic continuation *along the composed path*. The representative of a point of the universal covering $\text{cov}(D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}; x_j)$ is a pair (x, μ) , where x is a point of the punctured disk $D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}$, and $\mu \subset (D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\})$ is a path starting at the point x_j and ending at x . The path corresponding to the above mentioned two-step analytic continuation is the composition¹⁴ $\alpha = \mu \cdot \alpha_j$. Thus, to describe the dependence on $\mathbf{t}$ of the solution $Y_{\alpha_j}(x, \mathbf{t})$, which is normalized at the initial point x_0 by the condition

$$(6.18) \quad Y_{\alpha_j}(x, \mathbf{t})|_{x=x_0} = I,$$

in the neighborhood of the singular point t_j , we continue analytically this solution from the neighborhood $\mathcal{V}_{x_0}$ of the initial point to the punctured neighborhood $D_{t_j, \rho_j}(\mathbf{t}^0)$ along those paths α which reach first the point x_j (x_j does not depend on t_j and is common for all these paths), and then go, remaining within the punctured neighborhood $D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}$, to the destination points.

Definition 6.6. Such a family¹⁵ $\{Y_{\alpha_j}(x, \mathbf{t})\}_{\mathbf{t} \in \mathbf{D}(\mathbf{t}^0, \delta)}$ of analytic continuations (with respect to x) of the normalized by (6.18) solutions of the equation (6.1) from a neighborhood $\mathcal{V}_{x_0}$ of the initial point x_0 to the family $\{D_{t_j, \rho_j}(\mathbf{t}^0)\}_{t_j \in D_{t_j^0, \delta}}$ of the punctured disk neighborhoods of the points $t_j, t_j \in D_{t_j^0, \delta}$, is said to be **coherent** with respect to $\mathbf{t}$. The path α_j along which such continuations are carried out is said to be the **path determining the coherent family of analytic continuations**.

The solution

$$(6.19) \quad Y_j(x, \mathbf{t}) \cdot C_{\alpha_j}(\mathbf{t}) = H_j(x, \mathbf{t})(x - t_j)^{Q_j(\mathbf{t})} C_{\alpha_j}(\mathbf{t})$$

can be presented in another form – more suitable for our goal:

$$(6.20) \quad Y_{\alpha_j}(x, \mathbf{t}) = H_{\alpha_j}(x, \mathbf{t})(x - t_j)^{A_{\alpha_j}(\mathbf{t})},$$

where

$$(6.21) \quad \text{a). } A_{\alpha_j}(\mathbf{t}) = C_{\alpha_j}^{-1}(\mathbf{t}) Q_j(\mathbf{t}) C_{\alpha_j}(\mathbf{t}); \quad \text{b). } H_{\alpha_j}(x, \mathbf{t}) = H_j(x, \mathbf{t}) C_{\alpha_j}(\mathbf{t}).$$

¹⁴The path α starts at the point x_0 and ends at the point x . Traversing the path α , we traverse first the path α_j , and then the path μ . See Figure 5.b

¹⁵Indexed by the parameter $\mathbf{t} \in \mathbf{D}(\mathbf{t}^0, \delta)$

Of course, the relation (6.17) leads immediately to the relations (6.22)

$$\text{a). } A_{\alpha_j''}(\mathbf{t}) = (M_\gamma(\mathbf{t}))^{-1} A_{\alpha_j'}(\mathbf{t}) M_\gamma(\mathbf{t}), \quad \text{b). } H_{\alpha_j''}(x, \mathbf{t}) = H_{\alpha_j'}(x, \mathbf{t}) M_\gamma(\mathbf{t}),$$

where α_j' and α_j'' are two paths with the same endpoints x_0 and x_j , and M_γ is the monodromy matrix for the equation (6.1) corresponding to the loop γ , $\gamma = (\alpha_j')^{-1} \alpha_j''$.

Definition 6.7. The matrix functions $H_{\alpha_j}(x, \mathbf{t})$ and $(x - t_j)^{A_{\alpha_j}(\mathbf{t})}$ are said to be the *regular factor* and the *principal factor* of the representation (6.20) of the solution $Y_{\alpha_j}(x, \mathbf{t})$ near the point t_j .

Let us summarize the results, obtained above, as

Theorem 6.8. *Let the $k \times k$ matrix functions $Q_p(\mathbf{t})$, $p = 1, \dots, n$, which appear in the Fuchsian system (6.1) be holomorphic with respect to $\mathbf{t} = (t_1, \dots, t_n)$ for $\mathbf{t}$ from an open set $\mathcal{D}$, $\mathcal{D} \in \mathbb{C}_*^n$. Let $\mathbf{t}^0 = (t_1^0, \dots, t_n^0)$ be a point from $\mathcal{D}$. Assume that, for certain j , the matrix $Q_j(\mathbf{t}^0)$ is non-resonant, and let $\rho_j(\mathbf{t}^0)$ be defined according to (6.7)) (with $q = j$).*

Then there exists a number δ , $0 < \delta < \frac{1}{2}\rho_j(\mathbf{t}^0)$, such that, for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$, the solutions $\{Y_{\alpha_j}(x, \mathbf{t})\}_{\mathbf{t} \in D(\mathbf{t}^0, \delta)}$ of the holomorphic family of equation (6.1) which are distinguished by the properties:

- i. *Each of these solutions is normalized by (6.18) at the initial point x_0 ,*
- ii. *Each of these solution is continued analytically from a neighborhood of x_0 to the family $\{D_{t_j, \rho(\mathbf{t}^0)}\}_{t_j \in D_{t_j^0, \delta}}$ of the punctured disk neighborhoods of the points t_j , $t_j \in D_{t_j^0, \delta}$, and such analytic continuations are performed coherently with respect to $\mathbf{t} \in D(\mathbf{t}^0, \delta)$,*

are representable in $D_{t_j, \rho(\mathbf{t}^0)} \setminus \{t_j\}$ in the form (6.20), where:

- (1) *The matrix function $H_{\alpha_j}(x, \mathbf{t})$ — the regular factor of the representation (6.20) — is holomorphic together with its inverse $H_{\alpha_j}^{-1}(x, \mathbf{t})$ with respect to $x, \mathbf{t}$ for $x \in D_{t_j, \rho(\mathbf{t}^0)}$, $\mathbf{t} \in D(\mathbf{t}^0, \delta)$;*
- (2) *The exponent $A_{\alpha_j}(\mathbf{t})$ of the principal factor $(x - t_j)^{A_{\alpha_j}(\mathbf{t})}$ is a matrix function holomorphic with respect to $\mathbf{t}$ from the polydisk $\mathbf{t} \in D(\mathbf{t}^0, \delta)$;*
- (3) *The exponent $A_{\alpha_j}(\mathbf{t})$ is similar to the matrix $Q_j(\mathbf{t})$, (6.21.a), and the similarity matrix $C_{\alpha_j}(\mathbf{t})$ can be chosen¹⁶ to be holomorphic for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$.*
- (4) *By a different choice of the paths α_j' and α_j'' , along which the coherent analytic continuations are carried out, the matrices $A_{\alpha_j}(\mathbf{t})$ and $H_{\alpha_j}(x, \mathbf{t})$ are transformed according to the rule (6.22), where $M_\gamma(\mathbf{t})$*

¹⁶The similarity matrix is not unique: it can be multiplied by an arbitrary matrix commuting with $Q_j(\mathbf{t})$.

is the monodromy matrix corresponding to a loop γ , constructed¹⁷ from the paths α'_j and α''_j .

Let α_j be a path, separated (as in (6.11)) from the singular points $t_p, 1 \leq p \leq n$, and leading from the point x_0 to a point $x_j \in (\bigcap D_{t_j, \rho_j}(\mathbf{t}^0)) \setminus D_{t_j^0, \delta}$, where the intersection is taken over all $t_j \in D_{t_j^0, \delta}$. This path can be used for the coherent analytic continuations of the normalized by (6.18) solution to the neighborhoods of the point $t_j \in D_{t_j^0, \delta}$. The path α_j also generates a loop γ_{α_j} which goes around the disk neighborhood $D_{t_j^0, \delta}$. This loop does not depend on $\mathbf{t}$ and, for every $\mathbf{t} = (t_1, \dots, t_n), \mathbf{t} \in D(\mathbf{t}^0, \delta)$, represents an element of the fundamental group $\pi(\mathbb{C} \setminus \{t_1, \dots, t_n\}; x_0)$. (Compare to the Definition 5.12). The loop γ_{α_j} is the composition of the path α_j , of the circular arc μ , starting and ending at the point x_j , and of the path α_j^{-1} (the path α_j^{-1} is the path α_j , traversed in the “opposite direction”). In more detail, μ is the circular arc of the radius $|x_j - t_j^0|$, centered at the point t_j^0 , which starts at the point x_j , makes the full counterclockwise turn around the point t_j^0 , and ends at the same point x_j (or, better to say, at the point $t_j^0 + e^{2\pi i}(x_j - t_j^0)$). Since $x_j \in (\bigcap D_{t_j, \rho_j}(\mathbf{t}^0)) \setminus D_{t_j^0, \delta}$, $\mu \subset D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}$ for every $t_j \in D_{t_j^0, \delta}$. Since in $D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}$ the representation (6.20) holds, Theorem 5.18 can be formulated in the following “ $\mathbf{t}$ -dependent” version:

Theorem 6.9. *Assume that the matrix $Q_j(\mathbf{t}^0)$ is non-resonant. Let α_j be a path leading from the initial point x_0 to a punctured neighborhood of the singular point t_j^0 , and let γ_{α_j} be the loop generated by the path α_j (as described above).*

If $\mathbf{t} \in D(\mathbf{t}^0, \delta)$, and $\delta > 0$ is small enough, then the monodromy matrix $M_{\gamma_{\alpha_j}}(\mathbf{t})$ for the equation (6.1), corresponding to the loop γ_{α_j} , and the exponent $A_{\alpha_j}(\mathbf{t})$ of the principal factor of the representation (6.20) of its solution $Y_{\alpha_j}(x, \mathbf{t})$ are related by the equality

$$(6.23) \quad M_{\gamma_{\alpha_j}}(\mathbf{t}) = \exp\{2\pi i A_{\alpha_j}(\mathbf{t})\}.$$

The question arises, whether the monodromy matrix $M_{\gamma_{\alpha_j}}(\mathbf{t})$ determines the exponent $A_{\alpha_j}(\mathbf{t})$ uniquely? In particular, does the equality $A_{\alpha_j}(\mathbf{t}') = A_{\alpha_j}(\mathbf{t}'')$ follow from the equality $M_{\gamma_{\alpha_j}}(\mathbf{t}') = M_{\gamma_{\alpha_j}}(\mathbf{t}'')$, if $\mathbf{t}', \mathbf{t}'' \in D(\mathbf{t}^0, \delta)$, and $\delta \in$ is small enough?

The latter question is closely related to the description of the set of solutions of the matrix equation (with respect to A):

$$(6.24) \quad \exp\{2\pi i A\} = M,$$

where M is a given $k \times k$ matrix. A description of the set of solutions of (6.24) can be found in [Gant], Chapter VIII, §8. The invertibility of M is

¹⁷ Both paths α'_j and α''_j , have the common starting point x_0 . If their endpoints coincide, then the loop γ is just the composition $\gamma = (\alpha'_j)^{-1}\alpha''_j$. In the general case, in which these endpoints may be different, $\gamma = (\alpha'_j)^{-1}\nu\alpha''_j$, where ν is a path, connecting the endpoints.

the necessary and sufficient condition for the solvability of (6.24). Given $M, \det M \neq 0$, the set of solutions of (6.24) is always infinite. Our prime interest is in conditions under which a certain solution A_0 of the equation (6.24) is isolated. In general, non-isolated solutions may exist. For example, let $M = \text{diag}[1, 1]$, $A_0 = \text{diag}[0, 1]$, and let T be an arbitrary invertible matrix. Then $\exp\{2\pi i(T^{-1}A_0T)\} = M$. If T is non-diagonal matrix, then $T^{-1}A_0T \neq A_0$, but if T is close to I , then $T^{-1}A_0T$ is close to A_0 .

Lemma 6.10. *Given a non-resonant matrix A_0 , there exists $\varepsilon > 0$ such that from the conditions $\exp\{2\pi iA_0\} = \exp\{2\pi iA\}$ and $\|A - A_0\| < \varepsilon$ it follows that $A = A_0$.*

PROOF. Since A_0 is non-resonant, there exists an open (not necessarily connected) set $\mathcal{O}, \mathcal{O} \subset \mathbb{C}$ containing the spectrum of A_0 such that the mapping $\zeta \mapsto \exp\{2\pi i\zeta\}$ is univalent (schlicht) in $\mathcal{O}$. If ε is small enough, the set $\mathcal{O}$ also contains the spectra of all matrices A satisfying the conditions $\|A - A_0\| < \varepsilon$. The set $\mathcal{G} = \exp\{2\pi i\mathcal{O}\}$ is an open set containing all the spectra of the matrices $\exp\{2\pi iA\}$ with $\|A - A_0\| < \varepsilon$, in particular, the spectrum of the matrix $\exp\{2\pi iA_0\}$. If $\frac{1}{2\pi i}\text{Ln}w, \frac{1}{2\pi i}\text{Ln} : \mathcal{G} \mapsto \mathcal{O}$, is a function, inverse to the function $\exp\{2\pi i\zeta\} : \mathcal{O} \mapsto \mathcal{G}$, that is $\frac{1}{2\pi i}\text{Ln}(\exp\{2\pi i\zeta\}) \equiv \zeta \quad \forall \zeta \in \mathcal{O}$, then $\frac{1}{2\pi i}\text{Ln}(\exp\{2\pi iA\}) \equiv A$ for all matrices A whose spectra are contained in $\mathcal{O}$, in particular, $\forall A : \|A - A_0\| < \varepsilon$. Q.E.D.

Since the matrix $A_{\alpha_j}(\mathbf{t}^0)$ is similar to the matrix $Q_j(\mathbf{t}^0)$, the following result is the direct consequence of Lemma 6.10 and Theorem 6.9:

Theorem 6.11. *Let the matrix $Q_j(\mathbf{t}^0)$ be non-resonant, and let $A_{\alpha_j}(\mathbf{t})$ and $M_{\gamma_{\alpha_j}}(\mathbf{t})$ be the same as in the formulation of Theorem 6.9. If $\delta > 0$ is small enough, and if for some $\mathbf{t}', \mathbf{t}'' \in D(\mathbf{t}^0, \delta)$, the equalities $M_{\gamma_{\alpha_j}}(\mathbf{t}') = M_{\gamma_{\alpha_j}}(\mathbf{t}'')$ hold, then the equalities $A_{\alpha_j}(\mathbf{t}') = A_{\alpha_j}(\mathbf{t}'')$ hold as well. In particular, if $M_{\gamma_{\alpha_j}}(\mathbf{t}) \equiv M_{\gamma_{\alpha_j}}(\mathbf{t}^0)$ in $D(\mathbf{t}^0, \delta)$, then $A_{\alpha_j}(\mathbf{t}) \equiv A_{\alpha_j}(\mathbf{t}^0)$ in $D(\mathbf{t}^0, \delta)$ as well.*

Remark 6.12. If the matrix $Q_j(\mathbf{t}^0)$ is resonant, then, trying to formulate a “ $\mathbf{t}$ -dependent” version of Theorem 5.16, we should seek the representation of the solution $Y_{\alpha_j}(\mathbf{t})$ near $\mathbf{t}^0$ in the form

(6.25)

$$Y_{\alpha_j}(x, \mathbf{t}) = H_{\alpha_j}(x, \mathbf{t})(x - t_j)^{Z_{\alpha_j}(\mathbf{t})}(x - t_j)^{A_{\alpha_j}(\mathbf{t})}, \quad x \in \text{cov}(D_{t_j, \rho} \setminus \{t_j\}, x_j),$$

where the matrices $H_{\alpha_j}(x, \mathbf{t}), H_{\alpha_j}^{-1}(x, \mathbf{t})$ are holomorphic for $x \in D_{t_j, \rho}(\mathbf{t}^0)$ and $\mathbf{t} \in D(\mathbf{t}^0, \rho_j(\mathbf{t}^0))$, the matrices $Z_{\alpha_j}(\mathbf{t}), A_{\alpha_j}(\mathbf{t})$ are holomorphic for $\mathbf{t} \in D(\mathbf{t}^0, \rho_j(\mathbf{t}^0))$, and moreover the matrices $Z_{\alpha_j}(\mathbf{t})$ are diagonalizable, the eigenvalues of $Z_{\alpha_j}(\mathbf{t})$ are integer, the eigenvalues of $A_{\alpha_j}(\mathbf{t})$ and the eigenvalues of $Q_j(\mathbf{t})$ are related by equalities analogous to the equalities (5.29). However, if the matrix $Q_j(\mathbf{t}^0)$ is resonant, the representation of the form

(6.25), with holomorphic $Z_{\alpha_j}(\mathbf{t}), A_{\alpha_j}(\mathbf{t})$, does not exist in general. The reason is that it is possible for the matrix $Q_j(\mathbf{t}^0)$ to be resonant, while all the matrices $Q_j(\mathbf{t})$ for $\mathbf{t} \neq \mathbf{t}^0$ are non-resonant. (The property of resonance is unstable under small perturbations). According to Theorem 5.15, the solution $Y_{\alpha_j}(\mathbf{t})$ must be of the form (6.20) for $\mathbf{t} \neq \mathbf{t}^0$. According to Theorem 5.16, it must be of the form (6.25) for $\mathbf{t} = \mathbf{t}^0$, where in general $Z_{\alpha_j}(\mathbf{t}^0) \neq 0$. However, such a bifurcation is incompatible with the holomorphy of $Z_{\alpha_j}(\mathbf{t})$.

7. ISOPRINCIPAL AND ISOMONODROMIC DEFORMATIONS.

Consider a Fuchsian equation (6.1), where $k \times k$ matrices $Q_p(\mathbf{t}), 1 \leq p \leq n$, are holomorphic in $\mathbf{t}$ for $\mathbf{t}$ from a domain $\mathcal{D}, \mathcal{D} \subset \mathbb{C}_*^n$. The equation (6.1) is considered as a differential equation with respect to the variable x , and $\mathbf{t} = (t_1, \dots, t_n)$ is considered as a parameter. Recall that we use the terminology “the deformation of Fuchsian equation” for a Fuchsian equation depending on a parameter¹⁸.

Definition 7.1. Let (6.1) be a deformation of Fuchsian equation, where $k \times k$ matrices $Q_p(\mathbf{t}), 1 \leq p \leq n$, are holomorphic in a domain $\mathcal{D}, \mathcal{D} \subset \mathbb{C}_*^n$, and let $\mathbf{t}^0 = (t_1^0, \dots, t_n^0)$ be a point of $\mathcal{D}$. The deformation is said to be isomonodromic with respect to a distinguished point x_0 at the pole loci $\mathbf{t}^0$ if for every loop γ with the distinguished point $x_0, \gamma \subset \mathbb{C} \setminus \{t_1^0, \dots, t_n^0\}$, there exists $\delta > 0, \delta = \delta(\mathbf{t}^0, \gamma)$, such that the monodromy matrix $M_\gamma(\mathbf{t})$ of the deformation (6.1) along the loop γ does not depend on $\mathbf{t}$ in $D(\mathbf{t}^0, \delta)$: $M_\gamma(\mathbf{t}) \equiv M_\gamma(\mathbf{t}^0) \forall \mathbf{t} \in D(\mathbf{t}^0, \delta)$. (Of course, we assume that $\delta < \text{dist}(\mathbf{t}^0, \gamma)$, so that $\gamma \subset \mathbb{C} \setminus \{t_1, \dots, t_n\}$ for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$, and the monodromy matrix $M_\gamma(\mathbf{t})$ is well defined for such $\mathbf{t}$). The deformation is said to be isomonodromic with respect to x_0 in $\mathcal{D}$ if it is isomonodromic with respect to x_0 at every pole loci $\mathbf{t} \in \mathcal{D}$.

Remark 7.2. To check that the deformation (6.1) is isomonodromic at $\mathbf{t}^0$, there is no need to examine the monodromy matrix $M_\gamma(\mathbf{t})$ for all loops γ . It is enough to choose loops $\gamma_j, j = 1, \dots, n, \gamma_j \subset \mathbb{C} \setminus \{t_1^0, \dots, t_n^0\}$, generating¹⁹ the fundamental group $\pi(\mathbb{C} \setminus \{t_1^0, \dots, t_n^0\}, x_0)$, and to check the monodromy matrices $M_{\gamma_j}(\mathbf{t})$ do not depend on $\mathbf{t}$ for $\mathbf{t} \in D(\mathbf{t}^0, \delta), \delta > 0$ is small enough.

Remark 7.3. If γ is a loop with the distinguished point x_0 , and $t_j \notin \gamma, j = 1, \dots, n$, then the monodromy matrix $M_\gamma(\cdot)$ is holomorphic at the point $\mathbf{t} = (t_1, \dots, t_n)$. From the uniqueness theorem for holomorphic functions and from elementary topological considerations it follows that if the deformation (6.1) is isomonodromic at some $\mathbf{t} \in \mathcal{D}$, then it is isomonodromic at every $\mathbf{t} \in \mathcal{D}$.

¹⁸We consider only such deformations, where the loci $\mathbf{t} = (t_1, \dots, t_n)$ serve as a parameter, and the dependence on the parameter is holomorphic.

¹⁹If $\mathbf{t} \in D(\mathbf{t}^0, \delta)$ and $\delta > 0$ is small enough, then such loops γ_j satisfy the condition $\gamma_j \subset \mathbb{C} \setminus \{t_1, \dots, t_n\}$ and generate the fundamental group $\pi(\mathbb{C} \setminus \{t_1, \dots, t_n\}, x_0)$. See Figure 4.

Definition 7.4. Deformation (6.1) is said to be *isospectral*, if for every j , $j = 1, \dots, n$, the spectrum of the matrix $Q_j(\mathbf{t})$ does not depend on $\mathbf{t}$ in $\mathcal{D}$: $\text{spec } Q_j(\mathbf{t}') = \text{spec } Q_j(\mathbf{t}'')$ (the sets coincide “with multiplicities”) for every $\mathbf{t}', \mathbf{t}'' \in \mathcal{D}$.

Theorem 7.5. *If a deformation (6.1) is isomonodromic with respect to a distinguished point, then it is also isospectral.*

PROOF. For given j , the coefficients of the characteristic polynomials of the matrix $Q_j(\mathbf{t})$ are holomorphic with respect to $\mathbf{t}$ in $\mathcal{D}$. Therefore, it is enough to prove that for some $\delta > 0$, this polynomial does not depend on $\mathbf{t}$ for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$ or, what is the same, that the spectrum $\text{spec } Q_j(\mathbf{t})$ does not depend on $\mathbf{t}$ for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$. Let γ_j be a loop with the distinguished point x_0 which makes one turn counterclockwise around the singular point t_j^0 and makes no turns around other singular points t_p^0 , $p = 1, \dots, n, p \neq j$, as it was described in the formulation of Theorem 5.19. The monodromy matrix $M_{\gamma_j}(\mathbf{t})$ does not depend on $\mathbf{t}$ for $\mathbf{t}$ which are close to $\mathbf{t}^0$ (In general, $M_{\gamma_j}(\mathbf{t}) = M_{\gamma_j}(\mathbf{t}^0)$ if it is possible to pass from $\mathbf{t}^0$ to $\mathbf{t}$ in $\mathcal{D}$ so that no t_p crosses the loop γ_j). According to Theorem 5.19, the set $\exp\{\text{spec } Q_j(\mathbf{t})\}$ does not depend on $\mathbf{t}$ in $D(\mathbf{t}^0, \delta)$. However, on the one hand, the spectrum $\text{spec } Q_j(\mathbf{t})$ depends on $\mathbf{t}$ continuously, and on the other hand, the mapping $\zeta \mapsto \exp\{\zeta\}$ is locally univalent (schlicht). Therefore, the set $\text{spec } Q_j(\mathbf{t})$ does not depend on $\mathbf{t}$ in $D(\mathbf{t}^0, \delta)$ if δ is small enough. Q.E.D.

Let (6.1) be an isomonodromic with respect to x_0 deformation of Fuchsian equation, and let for a certain j , the matrix $Q_j(\mathbf{t}^0)$ be non-resonant²⁰. Let α_j , $\alpha_j \subset \mathbb{C} \setminus \{t_1^0, \dots, t_n^0\}$, be a path leading from the point x_0 to a small neighborhood of the point t_j^0 . According to Theorem 6.11, for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$, the solution $Y_{\alpha_j}(x, \mathbf{t})$, normalized at the distinguished point x_0 by the condition (6.12) and continued analytically to the neighborhood of t_j^0 , is representable near t_j in the form

$$(7.1) \quad Y_{\alpha_j}(x, \mathbf{t}) = H_{\alpha_j}(x, \mathbf{t})(x - t_j)^{A_{\alpha_j}}, \quad x \in D_{t_j, \rho}, \mathbf{t} \in D(\mathbf{t}^0, \delta),$$

where δ and ρ are some positive numbers, the *regular factor* $H(x, \mathbf{t})$ is holomorphic in $x, \mathbf{t}$ and invertible for $x \in D_{t_j, \rho}$, $\mathbf{t} \in D(\mathbf{t}^0, \delta)$, and the exponent A_{α_j} of the *principal factor* $(x - t_j)^{A_{\alpha_j}}$ *does not depend on $\mathbf{t}$* . Thus it is natural to give the following definition:

Definition 7.6. The deformation (6.1) is said to be *isoprincipal* with respect to a distinguished point x_0 in the narrow sense at the pole loci $\mathbf{t}^0 = (t_1^0, \dots, t_n^0)$, ($\mathbf{t}^0 \in \mathcal{D}$), if for every $j = 1, \dots, n$ and for some paths $\alpha_1, \dots, \alpha_n$ leading from x_0 to neighborhoods of the points $t_1^0, \dots, t_n^0$ respectively, the solution $Y_{\alpha_j}(x, \mathbf{t})$ normalized at x_0 by (6.12) is representable in the form (7.1) near

²⁰According to Theorem 7.5, the matrix $Q_j(\mathbf{t})$ is non-resonant for every $\mathbf{t} \in \mathcal{D}$.

t_j , where the matrices A_{α_j} do not depend on $\mathbf{t}$ for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$, $\delta > 0$ is small enough.

We emphasize that we do not require in this definition that the matrices $Q_j(\mathbf{t}^0)$ are non-resonant. Of course, in order for this definition to be correct, we must check that it does not depend on the choice of the paths α_j . This can be done easily. Below (Lemma 7.13) we prove this independence in a more general situation. Let us summarize the results in the form of two lemmas:

Lemma 7.7. *In the non-resonant case (i.e., if all the matrices $Q_j(\mathbf{t}^0)$ are non-resonant), every isomonodromic deformation is isoprincipal in the narrow sense as well.*

This lemma is the immediate consequence of Theorem 6.11. The converse is true always, without any assumption of non-resonance.

Lemma 7.8. *If the deformation (6.1) is isoprincipal (in the “narrow sense”) at some $\mathbf{t}^0$ (no assumptions on the matrices $Q_j(\mathbf{t}^0)$, $j = 1, \dots, n$ are made), then this deformation is isomonodromic as well.*

PROOF. If the solution $Y_{\alpha_j}(x, \mathbf{t})$ of (6.1) is representable in the form (7.1) with a constant A_{α_j} , then $M_{\gamma_{\alpha_j}}(\mathbf{t}) = \exp\{2\pi i A_{\alpha_j}\}$, where $M_{\gamma_{\alpha_j}}$ is the monodromy matrix along the loop γ_{α_j} generated by the path α_j . Therefore, $M_{\gamma_{\alpha_j}}$ does not depend on $\mathbf{t}$. Q.E.D.

Remark 7.9. If a deformation (6.1) is isomonodromic, but some of the matrices $Q_j(\mathbf{t})$ are not non-resonant, then this deformation may be not isoprincipal. As we shall see later, there is a rich class of deformations of Fuchsian equations all fundamental solutions $Y(x, \mathbf{t})$ of which are rational functions of x . Thus, such deformations are isomonodromic. (Their monodromies are trivial). However, most of these deformations are not isoprincipal.

From Lemmas 7.7 and 7.8 it follows that in the non-resonant case the classes of isomonodromic deformations and of isoprincipal (in the “narrow sense”) deformations coincide. This is a useful observation. On the one hand, it is the property of “isoprincipalness” that implies the Schlesinger system for the matrix functions $Q_p(\mathbf{t})$ appearing in the deformation. (We prove this implication below). On the other hand, there are methods, which are based on solving the so called Riemann-Hilbert problem, that allow to construct deformations of Fuchsian equations with the prescribed monodromy, but not with the prescribed principal factors of the solutions Y_{α_j} . In particular, such methods allow (under non-resonance condition and certain other restrictions) to construct directly isomonodromic (but not isoprincipal) deformations. The proof of the fact that the resulting $Q_j(\mathbf{t})$ satisfy the Schlesinger system uses essentially the equivalency of these two classes of deformations in the non-resonant case.

However, the above given “narrow” definition of the isoprincipalness, which is quite natural in the non-resonant case, seems to be a little bit

artificial and too restrictive in the general case. Indeed, in the general case the solutions $Y_{\alpha_j}(\mathbf{t})$ of the equation (6.1) are of the form (6.25), where exponents $Z_{\alpha_j}(\mathbf{t})$ and $A_{\alpha_j}(\mathbf{t})$ depend on $\mathbf{t}$ in general and possess some additional properties. In particular, the matrix $A_{\alpha_j}(\mathbf{t})$ is non-resonant, the matrix $Z_{\alpha_j}(\mathbf{t})$ is diagonalizable, with integer eigenvalues. (These exponents may be non-holomorphic, and even discontinuous functions of $\mathbf{t}$. See Remark 6.12). If these exponents are constant then we have good reason to call such a deformation isoprincipal. However, we have no need to go into detail. All that is important for us is that the principal factor $(x - t_j)^{Z_{\alpha_j}} \cdot (x - t_j)^{A_{\alpha_j}}$ depends on the difference $x - t_j$ only if the exponents Z_{α_j} and A_{α_j} do not depend on $\mathbf{t}$.

Definition 7.10. The deformation (6.1) is said to be isoprincipal with respect to a distinguished point x_0 (in the wide sense) at a pole loci $\mathbf{t}^0 = (t_1^0, \dots, t_n^0)$, ($\mathbf{t}^0 \in \mathcal{D}$), if for every $j = 1, \dots, n$ and for some paths $\alpha_1, \dots, \alpha_n$ leading from x_0 to neighborhoods of the points $t_1^0, \dots, t_n^0$ respectively, the solution $Y_{\alpha_j}(x, \mathbf{t})$ normalized at x_0 by (6.12) is representable in the form

$$(7.2) \quad Y_{\alpha_j}(x, \mathbf{t}) = H_{\alpha_j}(x, \mathbf{t}) E_{\alpha_j}(x - t_j), \quad x \in D_{t_j, \rho} \setminus \{t_j\}, \mathbf{t} \in \mathcal{D}(\mathbf{t}^0, \delta),$$

where the matrix functions $H_{\alpha_j}(x, \mathbf{t})$, $(H_{\alpha_j}(x, \mathbf{t}))^{-1}$ are holomorphic in $x, \mathbf{t}$ for $x \in D_{t_j, \rho}$, $\mathbf{t} \in \mathcal{D}(\mathbf{t}^0, \delta)$, (ρ and δ are some positive numbers), and $E_{\alpha_j}(\zeta)$, $j = 1, \dots, n$, are matrix functions possessing the properties:

- (1) $E_{\alpha_j}(\zeta)$ are holomorphic on the Riemann surface of logarithm²¹ and take invertible values there;
- (2) $E_{\alpha_j}(\zeta)$ do not depend on $\mathbf{t}$ and x .

Remark 7.11. Of course, to formulate Definition 7.10 more accurately, we have to describe the geometric configuration more accurately. For every j , we have to choose the point x_j according to (6.9), assuming that δ is small enough, in particular, (6.8) is satisfied. (The value $\rho_j(\mathbf{t}^0)$ is defined according to (6.7)). By such a choice of x_j , the condition (6.10) is satisfied. Then we have to choose the arguments $\arg(x_j - t_j^0)$ somehow, (6.14), and then $\arg(x_j - t_j)$ for $t_j \in D_{t_j^0, \delta}$ according to (6.15). A point of the universal covering $\text{cov}(D_{t_j, \rho_j(\mathbf{t}^0)} \setminus \{t_j\}, x_j)$ is a pair (x, μ) , where x is a point from the punctured disk $D_{t_j, \rho_j(\mathbf{t}^0)} \setminus \{t_j\}$, and μ is a path which starts at the point x_j and ends at the point x , $\mu \in D_{t_j, \rho_j(\mathbf{t}^0)} \setminus \{t_j\}$. The path μ , together with the choice of $\arg(x_j - t_j)$, determines the value of $\arg(x - t_j)$, and hence, the value of $E_{\alpha_j}(x - t_j)$. On the other hand, the pair $(x, \mu) \in \text{cov}(D_{t_j, \rho_j(\mathbf{t}^0)} \setminus \{t_j\}, x_j)$ determines the value of the analytic continuation Y_{α_j} along the path α_j at this point (x, μ) . By definition, the value $Y_{\alpha_j}((x, \mu), \mathbf{t})$ is the result of the analytic continuation of the solution normalized at x_0 by (6.18) along the composed path $\mu \cdot \alpha_j$ from a neighborhood of x_0 to the point x . Since both

²¹The Riemann surface of logarithm is the universal covering $\text{cov}(\mathbb{C} \setminus \{0\})$. Considered as functions defined in $\mathbb{C} \setminus \{0\}$, $E_{\alpha_j}(\zeta)$ are multivalued in general.

values, $Y_{\alpha_j}(x, \mathbf{t})$ and $E_{\alpha_j}(x - t_j)$ are determined for every choice of the path μ (these values may depend on μ), the value $H_{\alpha_j}(x, \mathbf{t}) \stackrel{\text{def}}{=} Y_{\alpha_j}(x, \mathbf{t})(E_{\alpha_j}(x - t_j))^{-1}$ is determined for $x \in D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}$, and it may depend on μ . The definition 7.10 means firstly, that the value $H_{\alpha_j}(x, \mathbf{t})$ is determined only by the point $x \in D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}$ itself, but does not depend on the path μ leading to this point, thus the function $H_{\alpha_j}(x, \mathbf{t})$ as a function of x is single-valued in the punctured disk $D_{t_j, \rho_j}(\mathbf{t}^0) \setminus \{t_j\}$. Secondly, this definition means that the singularity $x = t_j$ of the function $H_{\alpha_j}(x, \mathbf{t})$ is removable, that is there exists the limit $\lim_{x \rightarrow t_j, x \neq t_j} H_{\alpha_j}(x, \mathbf{t}) \stackrel{\text{def}}{=} H_{\alpha_j}(t_j, \mathbf{t})$. Thirdly, the definition 7.10 means that the value $H_{\alpha_j}(t_j, \mathbf{t})$ is an invertible matrix. (For $x \neq t_j$, the values $H_{\alpha_j}(t_j, \mathbf{t})$ are invertible, because both the values $Y_{\alpha_j}(x, \mathbf{t})$ and $E_{\alpha_j}(x - t_j)$ are invertible). So, the accurate definition of the isoprincipal deformation is rather bulky.

Below (Lemma 7.13) we prove that the notion of isoprincipal deformation is well defined, that is the property of the deformation to be isoprincipal at some point does not depend on the choice of the paths α_j .

Theorem 7.12. *If (6.1) is a deformation of Fuchsian equation which is isoprincipal with respect to a distinguished point in the wide sense (i.e, in the sense of Definition 7.10) at a pole loci $\mathbf{t}^0 = (t_1^0, \dots, t_n^0)$, ($\mathbf{t}^0 \in \mathcal{D}$), then the matrices $E_{\alpha_j}(\zeta)$, $j = 1, \dots, n$, representing the principle factors, must satisfy the monodromic relations of the form*

$$(7.3) \quad E_{\alpha_j}(\zeta e^{2\pi i}) = E_{\alpha_j}(\zeta) M_{\alpha_j}, \quad \forall \zeta \neq 0,$$

where M_{α_j} are constant (not depending on ζ) invertible matrices. For the monodromy matrix $M_{\gamma_{\alpha_j}}(\mathbf{t})$ of the equation (6.1) along the loop γ_{α_j} generated by the paths α_j , (see Figure 3), the equality

$$(7.4) \quad M_{\gamma_{\alpha_j}}(\mathbf{t}) = M_{\alpha_j}, \quad \mathbf{t} \in D(\mathbf{t}^0, \delta), \quad j = 1, \dots, n.$$

holds. In particular, this deformation is isomonodromic with respect to x_0 .

PROOF. For fixed $\mathbf{t} \in D(\mathbf{t}^0, \delta)$, and for $x \in D_{t_j, \rho} \setminus \{t_j\}$, consider two matrix functions of x : $Y_{\alpha_j}(x, \mathbf{t})$ and $Y_{\alpha_j}(t_j + (x - t_j)e^{2\pi i}, \mathbf{t})$. Here x and $t_j + (x - t_j)e^{2\pi i}$ are considered as points running over the universal covering $\text{cov}(D_{t_j, \rho} \setminus \{t_j\}, x_j)$ of the punctured disk $D_{t_j, \rho} \setminus \{t_j\}$ with the distinguished point x_j . These two points of the universal covering have the same projections in the complex plane. Therefore, the matrices $Y_{\alpha_j}(x, \mathbf{t})$ and $Y_{\alpha_j}(t_j + (x - t_j)e^{2\pi i}, \mathbf{t})$ can be considered as the values of two different solutions of the same linear differential equation at the same point. From $\det Y(x, \mathbf{t})|_{x=x_j} \neq 0$ (see (6.12)) it follows that $\det Y_{\alpha_j}(x, \mathbf{t}) \neq 0$, $\det Y_{\alpha_j}(t_j + (x - t_j)e^{2\pi i}, \mathbf{t}) \neq 0$. Hence, these solutions are “proportional” as functions of x :

$$(7.5) \quad Y_{\alpha_j}(t_j + (x - t_j)e^{2\pi i}, \mathbf{t}) = Y_{\alpha_j}(x, \mathbf{t}) M_{\alpha_j}(\mathbf{t}), \quad x \in D_{t_j, \rho} \setminus \{t_j\}, \mathbf{t} \in D(\mathbf{t}^0, \rho),$$

where $M_{\alpha_j}(\mathbf{t})$ are constant (with respect to x) non-degenerate matrices, which may depend on $\mathbf{t}$. Since the factor H_{α_j} is univalued ($H_{\alpha_j}(t_j + (x - t_j)e^{2\pi i}, \mathbf{t}) = H_{\alpha_j}(x, \mathbf{t})$), from (7.2) and (7.5) it follows that

$$(7.6) \quad E_{\alpha_j}((x - t_j)e^{2\pi i}) = E_{\alpha_j}(x - t_j)M_{\alpha_j}(\mathbf{t})$$

Since x is an arbitrary point of $D_{t_j, \rho} \setminus \{t_j\}$, $x - t_j$ is an arbitrary point of $\mathbb{C} \setminus \{0\}$. Since the function E_{α_j} does not depend on $\mathbf{t}$, the matrix M_{α_j} does not depend on $\mathbf{t}$ as well. Moreover, the monodromic relation (7.3) holds at least for $\zeta : 0 < |\zeta| < \rho$. However, the function E_{α_j} is holomorphic in the whole Riemann surface of logarithm. Therefore, the relation (7.3) holds for all $\zeta : |\zeta| > 0$.

The expression for the monodromy matrix $M_{\gamma_{\alpha_j}}(\mathbf{t})$ along the loop γ_{α_j} was already obtained (see (5.19)):

$$(7.7) \quad M_{\gamma_{\alpha_j}}(\mathbf{t}) = Y_{x_j, \alpha_j}(x_j, \mathbf{t})^{-1} \cdot Y_{x_j, \alpha_j}(t_j + (x_j - t_j)e^{2\pi i}, \mathbf{t}).$$

The equality (7.4) is the consequence of (7.5) and (7.7).

Q.E.D.

Let for a certain p , α_p be a path leading from the point x_0 to a neighborhood of the point t_p . Let α'_p be another path leading from x_0 to the same neighborhood. Let Y_{α_p} and $Y_{\alpha'_p}$ be the analytic continuations of the normalized at x_0 solution of (6.1) along the paths α_p and α'_p respectively. Then the solutions Y_{α_p} and $Y_{\alpha'_p}$ are related by the relation $Y_{\alpha'_p}(x, \mathbf{t}) = Y_{\alpha_p}(x, \mathbf{t}) \cdot M_\gamma(\mathbf{t})$, $x \in D_{t_p, \rho} \setminus \{t_j\}$, where $M_\gamma(\mathbf{t})$ is the monodromy matrix along the loop γ , constructed from the paths α_p and α'_p . (See footnote 17). If the deformation (6.1) is isoprincipal in the sense of the definition 7.10 by a certain choice of the paths $\alpha_j, j = 1, \dots, n$, then, according to Theorem 7.12, this deformation is isomonodromic. In particular, the monodromy matrix $M_\gamma(\mathbf{t})$ in (7.8) does not depend on $\mathbf{t}$: $M_\gamma(\mathbf{t}) = M_\gamma$, thus

$$(7.8) \quad Y_{\alpha'_p}(x, \mathbf{t}) = Y_{\alpha_p}(x, \mathbf{t}) \cdot M_\gamma, \quad x \in D_{t_p, \rho} \setminus \{t_j\},$$

Therefore, the solution $Y_{\alpha'_p}$ is representable in the form

$$(7.9) \quad Y_{\alpha'_p}(x, \mathbf{t}) = H_{\alpha'_p}(x, \mathbf{t}) \cdot E_{\alpha'_p}(x - t_j), \quad x \in D_{t_j, \rho} \setminus \{t_j\}, \mathbf{t} \in D(t_j, \rho),$$

with $H_{\alpha'_p}(x, \mathbf{t}) = H_{\alpha_p}(x, \mathbf{t})M_\gamma$, $E_{\alpha'_p}(\zeta) = M_\gamma^{-1}E_{\alpha_p}(\zeta)M_\gamma$. Since the matrices $E_{\alpha_p}(\zeta)$ and M_γ do not depend on $\mathbf{t}$, the matrix $E_{\alpha'_p}(\zeta)$ does not depend on $\mathbf{t}$ as well. So, we prove the following

Lemma 7.13. *The notion of the isoprincipal deformation at a point is well defined. If the deformation (6.1) is isoprincipal at the point $\mathbf{t}^0$ for some choice of the paths $\alpha_j, j = 1, \dots, n$ leading from x_0 to neighborhoods of the singularities t_j^0 , then this deformation is isoprincipal at this point for any other choice of such paths.*

Definition 7.14. The deformation (6.1), where the coefficients $Q_p(\mathbf{t})$, $p = 1, \dots, n$, are holomorphic for $\mathbf{t} \in \mathcal{D}$, $\mathcal{D}$ is an open set in $\mathbb{C}_*^n$, is said to be isoprincipal (in the wide sense) in $\mathcal{D}$ if it is isoprincipal at every point $\mathbf{t} \in \mathcal{D}$.

Theorem 7.15. *If the deformation (6.1) is isoprincipal with respect to x_0 at some $\mathbf{t}^0$, $\mathbf{t}^0 \in \mathcal{D}$, and the open set $\mathcal{D}$ is connected, then this deformation is isoprincipal with respect to x_0 in $\mathcal{D}$ (i.e., at every $\mathbf{t} \in \mathcal{D}$).*

We omit proof of this theorem.

8. ISOPRINCIPALNESS OF A DEFORMATION IMPLIES THE SCHLESINGER SYSTEM FOR ITS COEFFICIENTS

Lemma 8.1. *Let the deformation²²(6.1) be isoprincipal with respect to a distinguished point x_0 (in the wide sense) at some pole loci $\mathbf{t}^0$, $\mathbf{t}^0 \in \mathcal{D}$, and let $Y(x, \mathbf{t})$, $x \in \text{cov}(\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\})$, $\mathbf{t} \in \mathcal{D}$, be the solution of the differential equation (6.1) which is normalized by the condition (6.12) at the distinguished point x_0 .*

Then the equalities

$$(8.1) \quad \frac{\partial Y(x, \mathbf{t})}{\partial t_k} = Q_k(\mathbf{t}) \left(\frac{1}{x_0 - t_k} - \frac{1}{x - t_k} \right) Y(x, \mathbf{t}), \quad k = 1, \dots, n$$

are satisfied. In particular, if $x_0 = \infty$, then the equations (4.1b) are satisfied.

PROOF. The key to the proof are the (trivial, but nevertheless very useful) equalities

$$(8.2) \quad \frac{d}{dx} E_{\alpha_j}(x - t_j) = -\frac{\partial}{\partial t_j} E_{\alpha_j}(x - t_j), \quad j = 1, \dots, n,$$

which express the fact that the functions $E_{\alpha_j}(x - t_j)$ depend only on the difference of the arguments x and t_j . Expressing the logarithmic derivative $\frac{dY}{dx} Y^{-1}$ from the differential equation (6.1): $\frac{dY}{dx} Y^{-1} = \sum_p \frac{Q_p(\mathbf{t})}{x - t_p}$, we derive its behavior in a neighborhood of the singular point t_j :

$$(8.3) \quad \begin{aligned} \frac{dY}{dx}(x, \mathbf{t}) Y^{-1}(x, \mathbf{t}) &= \\ &= \frac{Q_j(\mathbf{t})}{x - t_j} + (\text{a function of } x, \text{ holomorphic with respect to } x), \end{aligned}$$

$$x \in D_{t_j, \rho}, \quad j = 1, \dots, n.$$

($D_{t_j, \rho} = \{z \in \mathbb{C} : |z - t_j| < \rho\}$, ρ is a positive number). We restrict our consideration to $\mathbf{t} \in D(\mathbf{t}^0, \delta)$, where $\delta > 0$ is small enough, so that the representations (7.2) of the solution $Y(x, \mathbf{t})$ hold in the appropriate

²²The condition $\sum_{1 \leq p \leq n} Q_p(\mathbf{t}) \equiv 0$, $\mathbf{t} \in \mathcal{D}$, is assumed.

neighborhoods $D_{t_j, \rho} \setminus \{t_j\}$ of the singular points t_j . Using the representation (7.2), we see that

$$(8.4) \quad \frac{dY_{\alpha_j}}{dx}(x, \mathbf{t}) Y_{\alpha_j}^{-1}(x, \mathbf{t}) = H_{\alpha_j}(x, \mathbf{t}) \left(\frac{d}{dx} E_{\alpha_j}(x - t_j) E_{\alpha_j}^{-1}(x - t_j) \right) H_{\alpha_j}^{-1}(x, \mathbf{t}) \\ + \left(\frac{d}{dx} H_{\alpha_j}(x, \mathbf{t}) \right) H_{\alpha_j}^{-1}(x, \mathbf{t}).$$

Since the function $H_{\alpha_j}(x, \mathbf{t})$ is holomorphic and invertible in an entire (non-punctured) neighborhood of the singular point t_j , comparing (8.3) and (8.4), we obtain

$$(8.5) \quad H_{\alpha_j}(x, \mathbf{t}) \left(\frac{d}{dx} E_{\alpha_j}(x - t_j) (E_{\alpha_j}(x - t_j))^{-1} \right) H_{\alpha_j}(x, \mathbf{t})^{-1} = \\ = \frac{Q_j}{x - t_j} + (\text{a function of } x, \text{ holomorphic with respect to } x), \\ x \in D_{t_j, \rho}, \quad j = 1, \dots, n.$$

Let us investigate the logarithmic derivatives $\frac{\partial Y}{\partial t_k}(x, \mathbf{t}) Y^{-1}(x, \mathbf{t})$, as functions of x . First of all we remark that for fixed $\mathbf{t}$, these functions are single valued and holomorphic with respect to x in the domain $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$. Indeed, let $\mathcal{O}$ be a simply connected open set in $\overline{\mathbb{C}}$ separated from the points $t_1^0, \dots, t_n^0$: $\text{dist}(\mathcal{O}, \{t_1^0, \dots, t_n^0\}) > 0$. Let $Y_1(x, \mathbf{t}), Y_2(x, \mathbf{t})$ be two branches of the solution $Y(x, \mathbf{t})$ in $\mathcal{O}$, that is the functions $Y_1(x, \mathbf{t}), Y_2(x, \mathbf{t})$ are obtained by the analytic continuation of the solution of (6.1) along two different paths γ_1, γ_2 leading from x_0 to $\mathcal{O}$. These branches $Y_1(x, \mathbf{t}), Y_2(x, \mathbf{t})$ must satisfy the relation $Y_2(x, \mathbf{t}) = Y_1(x, \mathbf{t}) M_\gamma(\mathbf{t})$, where $M_\gamma(\mathbf{t})$ is the monodromy matrix corresponding to the loop γ constructed from the paths γ_1 and γ_2 . In general, this monodromy matrix $M_\gamma(\mathbf{t})$ may depend on $\mathbf{t}$. However, according to Theorem 7.12, the considered deformation is isomonodromic. In particular, the matrix $M_\gamma(\mathbf{t})$ does not depend on $\mathbf{t}$, that is $Y_2(x, \mathbf{t}) = Y_1(x, \mathbf{t}) M_\gamma$, where M_γ does not depend on $\mathbf{t}$ and is invertible. (M_γ may depend on the choice of the branches Y_1 and Y_2 .) Differentiating the last equality with respect to t_k , we obtain $\frac{\partial Y_2}{\partial t_k} = \frac{\partial Y_1}{\partial t_k} M_\gamma$. From the last two equalities it follows that $\frac{\partial Y_1}{\partial t_k} Y_1^{-1} = \frac{\partial Y_2}{\partial t_k} Y_2^{-1}$, that is the logarithmic derivative $\frac{\partial Y}{\partial t_k} Y^{-1}$ is a single-valued holomorphic function of x in $\overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$. Using the representation (7.2) in a neighborhood of the singular point t_j , we see that for $j \neq k$

$$\frac{\partial Y}{\partial t_k}((x, \mathbf{t})) Y^{-1}((x, \mathbf{t})) = \frac{\partial H_{\alpha_j}(x, \mathbf{t})}{\partial t_k} H_{\alpha_j}^{-1}(x, \mathbf{t}), \quad x \in D_{t_j, \rho}.$$

(For $j \neq k$, the second factor $E_{\alpha_j}(x - t_j)$ in (7.2) does not depend on t_k .) Since the function $H_{\alpha_j}(x, \mathbf{t})$ is holomorphic with respect to $(x, \mathbf{t})$ for $x \in D_{t_j, \rho}$, $\mathbf{t} \in \mathcal{D}(\mathbf{t}^0, \delta)$, so is the function $\frac{\partial Y}{\partial t_k}((x, \mathbf{t}))Y^{-1}((x, \mathbf{t}))$ for $j \neq k$.

For $j = k$, we have to take into account the factor $E_{\alpha_j}(x - t_j)$:

$$\begin{aligned} \frac{\partial Y}{\partial t_k}((x, \mathbf{t}))Y^{-1}((x, \mathbf{t})) &= \\ &= H_{\alpha_k}(x, \mathbf{t}) \left(\frac{\partial}{\partial t_k} E_{\alpha_k}(x - t_k) E_{\alpha_k}^{-1}(x - t_k) \right) H_{\alpha_k}^{-1}(x, \mathbf{t}) \\ &\quad + \frac{\partial H_{\alpha_k}(x, \mathbf{t})}{\partial t_k} H_{\alpha_k}^{-1}(x, \mathbf{t}), \quad x \in D_{t_k, \rho}. \end{aligned}$$

The function $\frac{\partial H_{\alpha_k}(x, \mathbf{t})}{\partial t_k} H_{\alpha_k}^{-1}(x, \mathbf{t})$ is holomorphic for $x \in D_{t_j, \rho}$. Taking into account (8.2) and (8.5), we come to the conclusion:

$$(8.6) \quad \begin{aligned} \frac{\partial Y}{\partial t_k}((x, \mathbf{t}))Y^{-1}((x, \mathbf{t})) &= -\frac{Q_k}{x - t_k} + \\ &\quad + (\text{a function holomorphic with respect to } x), \quad x \in D_{t_k, \rho}. \end{aligned}$$

Thus, the function $\frac{\partial Y}{\partial t_k}((x, \mathbf{t}))Y^{-1}((x, \mathbf{t}))$, considered as a function of the variable x , is holomorphic in the entire extended complex plane $\overline{\mathbb{C}}$, except the point t_k . Near t_k this function behaves as it described in (8.6). Therefore,

$$\frac{\partial Y}{\partial t_k}((x, \mathbf{t}))Y^{-1}((x, \mathbf{t})) = -\frac{Q_k}{x - t_k} + C_k(\mathbf{t}),$$

where $C_k(\mathbf{t})$ does not depend on x . To determine $C_k(\mathbf{t})$, we use the normalizing condition (6.12). Since $Y(x_0, \mathbf{t}) \equiv I$, $\frac{\partial Y(x_0, \mathbf{t})}{\partial t_k} = 0$, i.e. $-\frac{Q_k(\mathbf{t})}{x - t_k} + C_k(\mathbf{t}) = 0$ at $x = x_0$. Thus, (8.1) holds. Q.E.D.

Theorem 8.2. *Let a $k \times k$ matrix functions $Q_j(\mathbf{t})$, $\mathbf{t} = (t_1, \dots, t_n)$, $j = 1, \dots, n$, are holomorphic for $\mathbf{t} \in \mathcal{D}$, where $\mathcal{D} \subset \mathbb{C}_*^n$ is an open connected set, and satisfying the condition*

$$\sum_{1 \leq j \leq n} Q_j(\mathbf{t}) = 0, \quad \mathbf{t} \in \mathcal{D}.$$

Assume that the deformation

$$\frac{dY}{dx} = \left(\sum_{1 \leq j \leq n} \frac{Q_j(\mathbf{t})}{x - t_j} \right) Y$$

is isoprincipal (in the wide sense) with respect to the distinguished point $x_0 = \infty$ at least for one pole loci $\mathbf{t}^0 \in \mathcal{D}$.

Then the matrix functions $Q_j(\mathbf{t})$, $j = 1, \dots, n$, satisfy the Schlesinger system (1.1) for all $\mathbf{t} \in \mathcal{D}$.

PROOF. Let $Y(x, \mathbf{t})$ be the solution of the differential equation (6.1) satisfying the normalizing condition $Y(x, \mathbf{t})_{x=\infty} = I$. According to Lemma 8.1, the equations (4.1b) are satisfied for $x \in \overline{\mathbb{C}} \setminus \{t_1, \dots, t_n\}$, $\mathbf{t} \in D(\mathbf{t}^0, \delta)$. The equation (4.1a) is the same as the equation (6.1). According to Theorem 4.1, the matrix functions $Q_j(\mathbf{t})$, $\mathbf{t} = (t_1, \dots, t_n)$, $j = 1, \dots, n$ satisfy the Schlesinger system (1.1) for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$. Since matrix functions in both left and right hand sides of (1.1) are holomorphic in $\mathcal{D}$ and coincide in $D(\mathbf{t}^0, \delta)$, they coincide everywhere in $\mathcal{D}$. Q.E.D.

9. CONSTRUCTION OF THE ISOPRINCIPAL DEFORMATIONS AS A TOOL FOR SOLVING THE SCHLESINGER SYSTEM.

Theorem 8.2 opens a way for constructing solutions of the Schlesinger system. The Cauchy problem for this system can be formulated as follows. Given $\mathbf{t}^0 \in \mathbb{C}_*^n$ and $k \times k$ matrices $Q_1^0, \dots, Q_n^0$, one needs to find matrix functions $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ of the variable $\mathbf{t}$ satisfying the Schlesinger system (1.1) and the initial condition

$$(9.1) \quad Q_j(\mathbf{t}^0) = Q_j^0, \quad j = 1, \dots, n.$$

If we succeed in constructing a family of Fuchsian equations

$$(9.2) \quad \frac{dY}{dx} = \left(\sum_{1 \leq j \leq n} \frac{Q_j(\mathbf{t})}{x - t_j} \right) Y,$$

enumerated by the pole loci $\mathbf{t} = (t_1, \dots, t_n)$, with holomorphic coefficients $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ (a holomorphic deformation) such that *this holomorphic deformation is isoprincipal with respect to $x_0 = \infty$* , then, according to Theorem 8.2, these coefficients will satisfy the Schlesinger system. In order for the initial conditions (9.1) to be satisfied, the family must contain the equation

$$(9.3) \quad \frac{dY}{dx} = \left(\sum_{1 \leq j \leq n} \frac{Q_j^0}{x - t_j^0} \right) Y,$$

which corresponds to the parameter value $\mathbf{t} = \mathbf{t}^0$.

If all the matrices $Q_1^0, \dots, Q_n^0$ are non-resonant (and hence, according to Theorems 7.12 and 7.5, all the matrices $Q_1(\mathbf{t}), \dots, Q_n(\mathbf{t})$ are non-resonant as well), this deformation is isoprincipal if, and only if, it is isomonodromic (Theorem 7.12 and Lemma 7.7). Therefore, if all the matrices $Q_1^0, \dots, Q_n^0$ are non-resonant, the Cauchy problem with initial conditions (9.1) is reduced to the problem of constructing the *isomonodromic* deformation (9.2) which in particular contains the equation (9.3). An approach to constructing such isomonodromic deformation lies in using *the Riemann-Hilbert problem*.

The Riemann-Hilbert problem can be formulated as follows.

Given $\mathbf{t} = (t_1, \dots, t_n) \in \mathbb{C}_*^n$, loops $\gamma_1(\mathbf{t}), \dots, \gamma_n(\mathbf{t})$ with the distinguished point x_0 that form the geometric configuration as it is plotted in Figure 1,

and matrices $M_1, \dots, M_n$ satisfying the condition²³ $M_{\pi(1)} \cdots M_{\pi(n)} = I$. One needs to construct the Fuchsian differential equation (9.2) with poles $t_1, \dots, t_n$ whose monodromy matrices take the prescribed values

$$(9.4) \quad M_{\gamma_j(t)} = M_j, \quad j = 1, \dots, n.$$

An approach for solving the Riemann-Hilbert problem was developed by J. Plemelj in 1908, and, a little later but independently, by G.D. Birkhoff (see [Plem1], [Plem2], [Birk]). The first step of this approach requires to solve a factorization problem for a matrix function on a closed contour. The contour passes through points $t_1, \dots, t_n$, and the matrix function is piecewise constant, constructed from the monodromy matrices $M_1, \dots, M_n$. This factorization problem can be reduced to a singular linear matrix integral equation on the contour. On the second step the solution of the factorization problem undergoes certain additional transformations in order to obtain the solution of the original Riemann-Hilbert problem. The second step is not completely painless: there are reefs here. In fact, not long ago A. A. Bolibruch has discovered examples of data for which the Riemann-Hilbert problem has no solution (see [AnBo], [Bol1], [Bol2], [Bol3]).

In order to apply the Riemann-Hilbert problem for the construction of the isomonodromic deformation, first we have to determine for the equation (9.3) (whose coefficients – matrices $Q_1^0, \dots, Q_n^0$ – are the initial values in the Cauchy problem for the Schlesinger system) the monodromy matrices $M_1^0, \dots, M_n^0$, corresponding to the loops $\gamma_1(t^0), \dots, \gamma_n(t^0)$. Then we have to solve the Riemann-Hilbert problem for the following data: an arbitrary point $t = (t_1, \dots, t_n) \in \mathbb{C}_*^n$, the appropriate loops $\gamma_1(t), \dots, \gamma_n(t)$, and the matrices $M_1^0, \dots, M_n^0$. As a result we shall obtain the matrices $Q_1(t), \dots, Q_n(t)$. Repeating this procedure for every t , (and, therefore, solving the infinite family of the Riemann-Hilbert problems) we construct matrix *functions* $Q_1(t), \dots, Q_n(t)$. If we succeed in performing this construction in such a way that these matrix functions are holomorphic then we obtain the desired isomonodromic deformation. Finally, if the initial data are non-resonant, we obtain the solution of the Cauchy problem for the Schlesinger system. Thus solving the Cauchy problem for a *non-linear* system is reduced to solving a family of *linear* problems.

However, solving the Schlesinger system in the non-resonant case is not the ultimate goal of the present manuscript. We would like to consider the case which is, in a certain sense, just the opposite. More precisely, we consider the initial values $Q_1^0, \dots, Q_n^0$ such that fundamental solutions of the Fuchsian system (9.3) are rational matrix functions of x in general position. Of course, a rational function is uni-valued in the complex plane, hence the monodromy in this case is trivial: $M_1 = \dots = M_n = I$. In this case the spectrum of each matrix Q_j^0 is either $\{0, 1\}$ or $\{0, -1\}$ and $\text{rank}(Q_j^0) =$

²³ $\{\pi_1, \dots, \pi_n\}$ is a permutation of the indices $\{1, \dots, n\}$ which is determined by the geometric configuration of the paths $\alpha_1, \dots, \alpha_n$. For the configuration plotted in Figure 1 the permutation is trivial: $\pi_k = k$.

1. In particular, all the matrices $Q_1^0, \dots, Q_n^0$ are resonant. Therefore, the considerations above, concerning the isomonodromic deformations, are not applicable in this case. Moreover, any family of Fuchsian systems with rational solutions is isomonodromic (the monodromies are trivial). Thus there are "too many" isomonodromic deformations here, and only one of them leads to the solution of the Schlesinger system. This is the isoprincipal deformation.

Let us recall Theorem 5.16 and, in particular, representation (5.28). If system (5.1) has a generic rational solution $Y(x)$ then in (5.28) we have $A_j = 0$, and hence $Y(x)$ is locally representable in a neighborhood of $x = t_j$ as

$$(9.5) \quad Y(x) = H_j(x)(x - t_j)^{Z_j},$$

with $H_j(x)$ holomorphic and invertible there, and

$$(9.6) \quad Z_j^2 = \pm Z_j.$$

Furthermore, under condition (9.6) the factor $(x - t_j)^{Z_j}$ has the following simple form:

$$(9.7) \quad (x - t_j)^{Z_j} = \begin{cases} I - Z_j + (x - t_j)Z_j, & \text{if } Z_j^2 = Z_j, \\ I + Z_j - \frac{1}{x - t_j}Z_j, & \text{if } Z_j^2 = -Z_j. \end{cases}$$

Thus the factorization (9.5) takes the form

$$(9.8) \quad Y(x) = H_j(x) \left(I - Z_j + (x - t_j)Z_j \right), \quad \text{if } Z_j^2 = Z_j,$$

$$(9.9) \quad Y(x) = H_j(x) \left(I + Z_j - \frac{1}{x - t_j}Z_j \right), \quad \text{if } Z_j^2 = -Z_j.$$

Factorizations of the form (9.8), (9.9) are traditional in the theory of rational matrix functions. The factors (9.7) are just the principal factors of the rational solution $Y(x)$ at the singular point t_j . If $Y(x, \mathbf{t})$ is a family of rational solutions for a family of Fuchsian systems which gives an isoprincipal deformation then for every $\mathbf{t} = (t_1, \dots, t_n)$ at each singular point $x = t_j$ $Y(x, \mathbf{t})$ must admit the factorization

$$(9.10) \quad Y(x, \mathbf{t}) = H_j(x, \mathbf{t}) \left(I - Z_j + (x - t_j)Z_j \right), \quad \text{if } Z_j^2 = Z_j,$$

$$(9.11) \quad Y(x, \mathbf{t}) = H_j(x, \mathbf{t}) \left(I + Z_j - \frac{1}{x - t_j}Z_j \right), \quad \text{if } Z_j^2 = -Z_j,$$

where Z_j do not depend on $\mathbf{t}$, and $H_j(x, \mathbf{t})$ are holomorphic and invertible at $x = t_j$. The realization theory for rational matrix functions allows to construct families of Fuchsian systems whose solutions $Y(x, \mathbf{t})$ admit the factorizations (9.10), (9.11). Development of the realization theory appropriate for this goal originated with L.A. Sakhnovich [Sakhn]. This topic was further developed in [GKRM]. In [Kats1] and [Kats2] this theory was adapted for applications concerning the Schlesinger system. The forthcoming second

part of the present manuscript will be dedicated to detailed construction of rational solutions of the Schlesinger system, based on the realization theory for rational matrix functions.

APPENDIX

A. THE REPRESENTATION OF A SOLUTION OF A FUCHSIAN EQUATION IN A NEIGHBORHOOD OF A SINGULAR POINT IN THE NON-RESONANT CASE.

PROOF OF THE PROPOSITION 6.3. Choose a positive δ so small that the closure of the polydisk $D(\mathbf{t}^0, \delta)$ is contained in the set $\mathcal{D}$, where the matrix functions $Q_p(\mathbf{t})$, $p = 1, \dots, n$, are defined and holomorphic. We also impose the condition $\delta < \frac{1}{2}\rho_j(\mathbf{t}^0)$. So, if $|t_j - t_j^0| < \delta$, and $|t_p - t_p^0| < \delta$, $p \neq j$, then $|t_j - t_p| > \rho_j(\mathbf{t}^0)$. Since the matrix $Q_j(\mathbf{t}^0)$ is non-resonant, choosing, if necessary, a smaller δ , we can also assume that the matrix $Q_j(\mathbf{t})$ is non-resonant for all $\mathbf{t} \in D(\mathbf{t}^0, \delta)$. Moreover, if $\delta > 0$ is small enough, then the differences of eigenvalues $\lambda_p(\mathbf{t}) - \lambda_q(\mathbf{t})$ of the matrix $Q_j(\mathbf{t})$ are separated from non-zero integers:

$$(A.1) \quad |\lambda_p(\mathbf{t}) - \lambda_q(\mathbf{t}) - m| \geq \varepsilon > 0 \quad \forall p, q = 1, \dots, k, \quad \forall m \in \mathbb{Z} \setminus \{0\}, \quad \forall \mathbf{t} \in D(\mathbf{t}^0, \delta),$$

where $\varepsilon > 0$ does not depend on $\mathbf{t} \in D(\mathbf{t}^0, \delta)$, $m \in \mathbb{Z} \setminus \{0\}$.

Changing variable

$$(A.2) \quad x \rightarrow x + t_j,$$

we reduce the differential equation (6.1) to the form

$$(A.3) \quad \frac{dZ(x, \mathbf{t})}{dx} = \left(\frac{Q_j(\mathbf{t})}{x} + \Phi(x, \mathbf{t}) \right) Z(x, \mathbf{t}),$$

where

$$(A.4) \quad Z(x, \mathbf{t}) = Y(x + t_j, \mathbf{t}), \quad \Phi(x, \mathbf{t}) = \sum_{\substack{1 \leq p \leq n \\ p \neq j}} \frac{Q_p(\mathbf{t})}{x + t_j - t_p}.$$

In view of the relation $|t_j - t_p| > \rho_j(\mathbf{t}^0)$, the function $\Phi(x, \mathbf{t})$ admits expansion of the form

$$(A.5) \quad \Phi(x, \mathbf{t}) = \sum_{0 \leq r < \infty} \Phi_r(\mathbf{t}) x^r, \quad x \in \mathbb{C}, \quad |x| < \rho_j(\mathbf{t}^0),$$

where $\Phi_r(\mathbf{t})$ are $k \times k$ matrix functions, which are holomorphic with respect to $\mathbf{t}$ for $\mathbf{t} \in D(\mathbf{t}^0, \delta)$ and admit the estimate

$$(A.6) \quad \|\Phi_r(\mathbf{t})\| \leq C_1 \cdot (\rho_j(\mathbf{t}^0))^{-r}, \quad \mathbf{t} \in D(\mathbf{t}^0, \delta), \quad 0 \leq r < \infty,$$

where $C_1 < \infty$ does not depend on $\mathbf{t}$ and r .

We seek a normalized solution of (A.3), corresponding to the singular point $x = 0$, in the form

$$(A.7) \quad Z(x) = \Psi(x, \mathbf{t}) x^{Q_j(\mathbf{t})},$$

where $\Psi(x, \mathbf{t})$ is power series in x :

$$(A.8) \quad \Psi(x, \mathbf{t}) = \sum_{0 \leq r < \infty} \Psi_r(\mathbf{t}) x^r, \quad \text{with} \quad \Psi_0(\mathbf{t}) = I.$$

Substituting the expressions (A.7) - (A.8) and (A.5) into (A.3) and using the differentiation rule

$$(A.9) \quad \frac{d}{dx} x^{Q_j(\mathbf{t})} = \frac{Q_j(\mathbf{t})}{x} x^{Q_j(\mathbf{t})},$$

we obtain for $r = 0, 1, 2, \dots$ the equalities

$$(A.10) \quad (r+1)\Psi_{r+1}(\mathbf{t}) + \Psi_{r+1}(\mathbf{t})Q_j(\mathbf{t}) - Q_j(\mathbf{t})\Psi_{r+1}(\mathbf{t}) = \sum_{\substack{l+m=r \\ l \geq 0, m \geq 0}} \Phi_l(\mathbf{t})\Psi_m(\mathbf{t}),$$

relating the given sequence $\{\Phi_r(\mathbf{t})\}_{0 \leq r < \infty}$ and the sequence $\{\Psi_r(\mathbf{t})\}_{0 \leq r < \infty}$, which has to be found. The relations (A.10) express the fact that (A.7) - (A.8) is a *formal solution* of the differential equation (6.1). If the series (A.8) converge in a disk $\{x \in \mathbb{C} : |x| < \rho\}$, with $\rho > 0$, then (A.7), where $\Psi(x, \mathbf{t})$ is the sum of these series, is an actual solution of (6.1) in the punctured disk $\{x \in \mathbb{C} : |x| < \rho\} \setminus \{0\}$.

Since the constant term $\Psi_0(\mathbf{t})$ is given: $\Psi_0(\mathbf{t}) = I$, the equations (A.10) can be considered as a recursive system for successive determining of $\Psi_1(\mathbf{t})$, $\Psi_2(\mathbf{t})$, etc. The r -th equation of the system (A.10) is of the form

$$(A.11) \quad \lambda X - [Q, X] = Y,$$

where $[Q, X] = QX - XQ$ — the commutator of the matrices Q and X . In (A.10),

$$(A.12) \quad \lambda = r+1, \quad Q = Q_j(\mathbf{t}), \quad X = \Psi_{r+1}(\mathbf{t}), \quad Y = \sum_{\substack{l+m=r \\ l \geq 0, m \geq 0}} \Phi_l(\mathbf{t})\Psi_m(\mathbf{t}).$$

Now we interrupt the proof of Proposition 6.3 to discuss the matrix equation (A.11).

Lemma A.1. *Let $\mathfrak{M}_k$ be the set of all $k \times k$ matrices with complex entries. Given a matrix $Q \in \mathfrak{M}_k$, let us associate with this matrix the operator ad_Q :*

$$(A.13) \quad \text{ad}_Q X = QX - XQ, \quad \text{ad}_Q : \mathfrak{M}_k \rightarrow \mathfrak{M}_k.$$

Let $\lambda_1(Q), \dots, \lambda_k(Q)$ be the set of all eigenvalues of the matrix Q (enumerated with multiplicities). Then the set of all eigenvalues (enumerated with multiplicities) of the operator ad_Q is the set of all the differences $\{\lambda_p(Q) - \lambda_q(Q)\}_{1 \leq p, q \leq k}$.

PROOF. Assume first that all eigenvalues of the matrix Q are pairwise different. Let u_p and v_q be, respectively, eigenvector columns and eigenvector rows of the matrix Q :

$$Qu_p = \lambda_p u_p, \quad v_q Q = \lambda_q v_q, \quad 1 \leq p, q \leq k.$$

Then the matrices $X_{p,q} \stackrel{\text{def}}{=} u_p \otimes v_q$ are eigenvectors of the operator ad_Q :

$$\text{ad}_Q X_{p,q} = (\lambda_p - \lambda_q) X_{p,q}, \quad 1 \leq p, q \leq k.$$

Thus, the numbers $\lambda_p(Q) - \lambda_q(Q)$ are eigenvalues of the operator ad_Q . Since the eigenvalues of the matrix Q are pairwise different, the vectors $\{u_p\}_{1 \leq p \leq k}$ and $\{v_q\}_{1 \leq q \leq k}$ form bases in the spaces of all k -vector columns and k -vector rows respectively. Since dimension of the space $\mathfrak{M}_k$ is equal to k^2 , the set of matrices $\{X_{p,q}\}_{1 \leq p,q \leq k}$ is a basis of the space $\mathfrak{M}_k$. Therefore, the set $\{\lambda_p(Q) - \lambda_q(Q)\}_{1 \leq p,q \leq k}$ is the set of all eigenvalues of the operator ad_Q (enumerated with multiplicities). This result can be extended to general matrices Q , i.e. to matrices whose eigenvalues are not necessarily pairwise different. Such an extension can be done using the approximation reasoning. Q.E.D.

From the above result it follows that if λ satisfies the condition $\lambda \neq \lambda_p(Q) - \lambda_q(Q)$,

$1 \leq p, q \leq k$, then the equation (Q, X, Y are $k \times k$ matrices)

$$(A.14) \quad \lambda X - \text{ad}_Q X = Y$$

is solvable with respect to X for every Y . However, for what follows one needs not only the solvability of this equation but also an estimate for its solution.

Lemma A.2. *Let $\mathcal{L}$ be a K -dimensional vector space equipped with a norm $\|\cdot\|_{\mathcal{L}}$, and let $\mathcal{A}$ be a linear operator acting in $\mathcal{L}$. Assume that the eigenvalues $\lambda_m(\mathcal{A})$ of the operator $\mathcal{A}$ are separated from zero:*

$$(A.15) \quad |\lambda_m(\mathcal{A})| \geq \varepsilon, \quad 1 \leq m \leq K.$$

Then the inverse operator $\mathcal{A}^{-1}$ admits the estimate from above:

$$(A.16) \quad \|\mathcal{A}^{-1}\| \leq 2^K \|\mathcal{A}\|^{K-1} \varepsilon^{-K},$$

where $\|\mathcal{A}\| = \sup \| \mathcal{A}x \|_{\mathcal{L}}, \|\mathcal{A}^{-1}\| = \sup \| \mathcal{A}^{-1}x \|_{\mathcal{L}}$, and sup's are taken over all $x \in \mathcal{L}$ such that $\|x\|_{\mathcal{L}} \leq 1$.

PROOF. Let $\chi(\zeta) = \det(\zeta \mathcal{I} - \mathcal{A})$ be the characteristic polynomial of the operator $\mathcal{A}$,

$$\chi(\zeta) = \zeta^K - \sigma_1 \zeta^{K-1} + \dots + (-1)^{K-1} \sigma_{K-1} \zeta + (-1)^K \sigma_K,$$

where $\sigma_l, 1 \leq l \leq K$, denotes the l -th elementary symmetric polynomial of in $\lambda_1(\mathcal{A}), \dots,$

$\lambda_K(\mathcal{A})$. According to Hamilton-Cayley theorem, $\chi(\mathcal{A}) = 0$, hence

$$(A.17) \quad \mathcal{A}^{-1} = (-1)^{K-1} \frac{1}{\sigma_K} \mathcal{A}^{K-1} + (-1)^{K-2} \frac{\sigma_1}{\sigma_K} \mathcal{A}^{K-2} + \dots + \frac{\sigma_{K-1}}{\sigma_K} \mathcal{I}.$$

From (A.17) it follows that

$$\|\mathcal{A}^{-1}\| \leq \left| \frac{\sigma_0}{\sigma_K} \right| \|\mathcal{A}\|^{K-1} + \left| \frac{\sigma_1}{\sigma_K} \right| \|\mathcal{A}\|^{K-2} + \dots + \left| \frac{\sigma_{K-1}}{\sigma_K} \right|. \quad (\sigma_0 \stackrel{\text{def}}{=} 1)$$

It is clear that the fractions $\frac{\sigma_l}{\sigma_K}$ are the elementary symmetric polynomials in the inverse eigenvalues $\frac{1}{\lambda_1(\mathcal{A})}, \dots, \frac{1}{\lambda_K(\mathcal{A})}$. From (A.15) we conclude that

$$\left| \frac{\sigma_l}{\sigma_K} \right| \leq C_K^l \varepsilon^{l-K}, \quad \text{where } C_K^l \text{ are binomial coefficients: } C_K^l = \frac{K!}{l!(K-l)!}.$$

Therefore,

$$\|\mathcal{A}^{-1}\| \leq C_K^0 \varepsilon^{-K} \|\mathcal{A}\|^{K-1} + C_K^1 \varepsilon^{1-K} \|\mathcal{A}\|^{K-2} + \dots + C_K^{K-1} \varepsilon^{-1} \|\mathcal{A}\|^0.$$

Using the binomial formula, we rewrite the previous inequality as

$$(A.18) \quad \|\mathcal{A}^{-1}\| \leq \|\mathcal{A}\|^{-1} \left((1 + \varepsilon^{-1} \|\mathcal{A}\|)^K - 1 \right),$$

Since $|\lambda_m(\mathcal{A})| \leq \|\mathcal{A}\|$ for any eigenvalue $\lambda_m(\mathcal{A})$, inequality (A.15) implies that $1 \leq \varepsilon^{-1} \|\mathcal{A}\|$, and hence $1 + \varepsilon^{-1} \|\mathcal{A}\| \leq 2\varepsilon^{-1} \|\mathcal{A}\|$. Weakening the inequality (A.18), we come to the more rough, but more simple inequality (A.16). Q.E.D.

Lemma A.3. *Let Q be a $k \times k$ matrix, with eigenvalues $\lambda_1, \dots, \lambda_k$, enumerated with multiplicities. Let λ be a complex number, and let for some $\varepsilon > 0$ the inequalities*

$$(A.19) \quad |\lambda - (\lambda_p - \lambda_q)| > \varepsilon, \quad 1 \leq p, q \leq k,$$

hold true. Then for every Y , the equation (A.14) (or, what is the same, the equation (A.11)) is solvable with respect to X , and the estimate

$$(A.20) \quad \|X\| \leq 2^{k^2} \varepsilon^{-k^2} (|\lambda| + 2\|Q\|)^{k^2-1} \|Y\|,$$

holds, where $\|\cdot\|$ is an arbitrary norm on $\mathfrak{M}_k$ with the property: $\|AB\| \leq \|A\| \|B\|$ for every $A, B \in \mathfrak{M}_k$.

PROOF. We use Lemma A.2 for the operator $\mathcal{A} = \lambda \mathcal{I} - \text{ad}_Q$, acting on the space $\mathfrak{M}_k$. It is clear that $\|\mathcal{A}\| \leq |\lambda| + 2\|Q\|$. According to Lemma A.1, the set $\{\lambda - (\lambda_p - \lambda_q)\}_{1 \leq p, q \leq k}$ is the set of all eigenvalues of $\mathcal{A}$ (enumerated with multiplicities). The inequality (A.19) provides an estimate of these eigenvalues from below. Applying Lemma A.1 to this operator $\mathcal{A}$ (and taking into account that $\dim \mathfrak{M}_k = k^2$), we obtain the estimate (A.20). Q.E.D.

The following lemma gives an estimate, which is more precise than (A.20) for large λ .

Lemma A.4. *Let Q be a $k \times k$ matrix, with eigenvalues $\lambda_1, \dots, \lambda_k$, enumerated with multiplicities. Let λ be a complex number which is different from the numbers $\lambda_p - \lambda_q$, $1 \leq p, q \leq k$, and let the estimates for $\|Q\|$ from above and for $|\lambda - (\lambda_p - \lambda_q)|$ from below be given:*

$$(A.21) \quad |\lambda - (\lambda_p - \lambda_q)| \geq \varepsilon, \quad 1 \leq p, q \leq k, \quad \|Q\| \leq \mu, \quad (\varepsilon > 0, \mu < \infty).$$

Then for every $Y \in \mathfrak{M}_k$, the equation (A.14) (or, what is the same, the equation (A.11)) is solvable with respect to $X \in \mathfrak{M}_k$, and the estimate

$$(A.22) \quad \|X\| \leq \frac{C(\varepsilon, \mu, k)}{1 + |\lambda|} \|Y\|,$$

holds, where $\|\cdot\|$ is an arbitrary norm on $\mathfrak{M}_k$ with the property: $\|AB\| \leq \|A\|\|B\|$ for every $A, B \in \mathfrak{M}_k$, and $C(\varepsilon, \mu, k) < \infty$ is a constant depending only on ε, μ and k .

PROOF. If $\lambda X - \text{ad}_Q X = Y$, then $|\lambda|\|X\| \leq \|\text{ad}_Q X\| + \|Y\| \leq 2\|Q\|\|X\| + \|Y\| \leq 2\mu\|X\| + \|Y\|$. Thus,

$$(A.23) \quad \|X\| \leq \frac{1}{|\lambda| - 2\mu} \|Y\|, \quad \text{for } |\lambda| > 2\mu.$$

On the other hand, if we assume that $|\lambda| \leq 3\mu$ and replace in the inequality (A.20) $|\lambda|$ and $\|Q\|$ by the larger values 3μ and μ respectively, we come to the inequality

$$(A.24) \quad \|X\| \leq 10^{k^2} \varepsilon^{-k^2} \mu^{k^2-1} \|Y\|, \quad \text{for } |\lambda| \leq 3\mu.$$

Unifying the inequalities (A.23) and (A.24), we come to the inequality (A.22) with a suitable constant $C(\varepsilon, \mu, k)$ (which, of course, can be found explicitly). Q.E.D.

Now, after we have investigated the matrix equation (A.11), we resume the proof of Proposition 6.3. We apply Lemma A.3 to the equality (A.10) considered as an equation of the form (A.11), with X, Y, Q, λ defined in (A.12). The number $\max_{t \in D(t^0)} \|Q_j(t)\|$ serves as μ , the number ε is taken from

(A.1). Substituting these X, λ and the estimate

$$(A.25) \quad \|Y\| \leq \sum_{0 \leq m \leq r} \|\Phi_{r-m}(t)\| \|\Psi_m(t)\|$$

for $\|Y\|$ into (A.22), we obtain, for $t \in D(t^0, \delta)$, $r = 0, 1, 2, \dots$ the estimate

$$(A.26) \quad \|\Psi_{r+1}(t)\| \leq \frac{C(\varepsilon, \mu, k)}{r+2} \sum_{0 \leq m \leq r} \|\Phi_{r-m}(t)\| \|\Psi_m(t)\|.$$

Substituting the upper estimate (A.6) for $\|\Phi_{r-m}(t)\|$ into (A.26), we obtain:

$$(A.27) \quad \|\Psi_{r+1}(t)\| \rho^{r+1} \leq \frac{C_2}{r+2} \sum_{0 \leq m \leq r} \|\Psi_m(t)\| \rho^m, \quad \forall t \in D(t^0, \delta), \quad r = 0, 1, 2, \dots,$$

where

$$(A.28) \quad C_2 = \rho \cdot C_1 \cdot C(\varepsilon, \mu, k),$$

$\rho = \rho_j(\mathbf{t}^0)$ is defined in (6.7), C_1 and $C(\varepsilon, \mu, k)$ are the values from (A.6) and (A.22) respectively. It is important to note that the value $C_2 < \infty$ does not depend on r and on $\mathbf{t} \in D(\mathbf{t}^0)$.

Lemma A.5. *Let $\{a_r\}_{0 \leq r < \infty}$ be a sequence of non-negative numbers, satisfying the conditions*

$$(A.29) \quad a_{r+1} \leq \frac{d}{r+2} \sum_{0 \leq m \leq r} a_m, \quad r = 0, 1, 2, \dots; \quad a_0 = 1,$$

where $d, 0 < d < \infty$, does not depend on r .

Then the sequence $\{a_r\}_{0 \leq r < \infty}$ admits the upper estimate

$$(A.30) \quad a_r \leq (r+1)^{d-1}, \quad r = 0, 1, 2, \dots$$

PROOF. For $r = 0$, the assertion (A.30) is true: $a_0 = 1$. Assume that the inequalities (A.30) hold for $r = 0, 1, \dots, m$. Then, according to (A.30), the inequality

$$(A.31) \quad a_{m+1} \leq \frac{d}{m+2} \sum_{0 \leq r \leq m} (r+1)^{d-1}$$

holds. If $d \geq 1$, then

$$(r+1)^{d-1} \leq \int_{r+1}^{r+2} t^{d-1} dt, \quad \sum_{0 \leq r \leq m} (r+1)^{d-1} \leq \int_1^{m+2} t^{d-1} dt \leq \frac{(m+2)^d}{d},$$

and the inequality (A.30) holds for $r = m+1$. If $0 < d < 1$, then

$$(r+1)^{d-1} \leq \int_r^{r+1} t^{d-1} dt, \quad \sum_{0 \leq r \leq m} (r+1)^{d-1} \leq \int_0^{m+1} t^{d-1} dt \leq \frac{(m+2)^d}{d},$$

and the inequality (A.30) holds for $r = m+1$ as well. By induction, the inequality (A.30) holds for all r . Q.E.D.

Applying Lemma A.5 to (A.27), with $a_r = \|\Psi_r(\mathbf{t})\| \rho^r$, we obtain the estimate

$$(A.32) \quad \|\Psi_r(\mathbf{t})\| \leq (r+1)^d \rho^{-r}, \quad r = 0, 1, 2, \dots, \quad \forall \mathbf{t} \in D(\mathbf{t}^0, \delta),$$

where $\rho = \rho_j(\mathbf{t}^0)$ is defined in (6.7), $d = C_2$ from (A.28), $d < \infty$ does not depend on r and on $\mathbf{t} \in D(\mathbf{t}^0, \delta)$.

From the estimate (A.32) it follows, that the power series (A.8) converge locally uniformly in the open circle $\{x : |x| < \rho\}$, and the convergence is uniform with respect to the parameter $\mathbf{t} \in D(\mathbf{t}^0, \delta)$. Therefore, the sum $\Psi(x, \mathbf{t})$ of these power series is a function, holomorphic with respect to $x, \mathbf{t}$ for $\{x : |x| < \rho_j(\mathbf{t}^0)\}, \mathbf{t} \in D(\mathbf{t}^0, \delta)$. Let us prove that the function $\Psi^{-1}(x, \mathbf{t})$

is holomorphic for these ²⁴ $x, \mathbf{t}$ as well. Since $\Psi_0(\mathbf{t}) = I$, we can consider the *formal* power series

$$(A.33) \quad \Psi^{\text{inv}}(x, \mathbf{t}) = \sum_{0 \leq r < \infty} \Psi_r^{\text{inv}}(\mathbf{t}) x^r,$$

which are *inverse* to the series (A.8), i.e.

$$(A.34) \quad \left(\sum_{0 \leq r < \infty} \Psi_r(\mathbf{t}) x^r \right) \cdot \left(\sum_{0 \leq r < \infty} \Psi_r^{\text{inv}}(\mathbf{t}) x^r \right) = \left(\sum_{0 \leq r < \infty} \Psi_r^{\text{inv}}(\mathbf{t}) x^r \right) \cdot \left(\sum_{0 \leq r < \infty} \Psi_r(\mathbf{t}) x^r \right) = I.$$

It is clear that

$$(A.35) \quad \Psi_0^{\text{inv}}(\mathbf{t}) = I.$$

The expressions (A.3), (A.7), (A.8) and (A.34) imply that the product

$$(A.36) \quad Z^{\text{inv}}(x, \mathbf{t}) \stackrel{\text{def}}{=} x^{-Q_j(\mathbf{t})} \Psi^{\text{inv}}(x, \mathbf{t})$$

is a formal solution of the differential equation

$$(A.37) \quad \frac{dZ^{\text{inv}}}{dx} = -Z^{\text{inv}} \left(\frac{Q_j(\mathbf{t})}{x} + \Phi(x, \mathbf{t}) \right),$$

As before (see (A.10)), substituting the expression (A.36) into (A.37), we obtain the infinite recursive system

$$(A.38) \quad (r+1)\Psi_{r+1}^{\text{inv}}(\mathbf{t}) + \Psi_{r+1}^{\text{inv}}(\mathbf{t})Q_j(\mathbf{t}) - Q_j(\mathbf{t})\Psi_{r+1}^{\text{inv}}(\mathbf{t}) = - \sum_{\substack{l+m=r \\ l \geq 0, m \geq 0}} \Psi_m^{\text{inv}}(\mathbf{t})\Phi_l(\mathbf{t}).$$

Like the r -th equation of the system (A.10), the r -th equation of the system (A.38) is of the form (A.11), with *the same* λ and Q as in (A.10): $\lambda = r+1, Q = Q_j(\mathbf{t})$. The only difference is that the right-hand side of (A.11) is $Y = \sum \Phi_l(\mathbf{t})\Psi_m(\mathbf{t})$ for (A.10), and $Y = -\sum \Psi_m^{\text{inv}}(\mathbf{t})\Phi_l(\mathbf{t})$ for (A.38). Because λ and Q for both matrix equations, corresponding to the systems (A.10) and (A.38), are the same, the estimates (A.22) for both equations hold with the same constant $C(\varepsilon, \mu, k)$ (Of course, ε, μ, k are the same). Instead of the estimate (A.25), the estimate

$$(A.39) \quad \|Y\| \leq \sum_{0 \leq m \leq r} \|\Phi_{r-m}(\mathbf{t})\| \|\Psi_m^{\text{inv}}(\mathbf{t})\|$$

²⁴Of course, in view of the condition $\Psi_0(\mathbf{t}) = I$, the function $\Psi^{-1}(x, \mathbf{t})$ is holomorphic for x which are small enough. However, it is very easy to prove the holomorphy of the function Ψ^{-1} in the whole disc $\{x : |x| < \rho\}$. Such a proof can be done "for free", using the previously established estimates.

for the right-hand side of (A.11) should be used now. Finally, we come to the recursive system of inequalities

$$(A.40) \quad \|\Psi_{r+1}^{\text{inv}}(\mathbf{t})\| \leq \frac{C(\varepsilon, \mu, k)}{r+2} \sum_{0 \leq m \leq r} \|\Phi_{r-m}(\mathbf{t})\| \|\Psi_m^{\text{inv}}(\mathbf{t})\|, \quad \forall \mathbf{t} \in D(\mathbf{t}^0, \delta).$$

with respect to $\|\Psi_m^{\text{inv}}(\mathbf{t})\|$. However, the latter system of inequalities coincides with the system (A.26) of inequalities with respect to $\|\Psi_m(\mathbf{t})\|$. Since $\|\Psi_0^{\text{inv}}(\mathbf{t})\| = \|\Psi_0(\mathbf{t})\| = \|I\|$, the estimates for $\|\Psi_m(\mathbf{t})\|$ and $\|\Psi_m^{\text{inv}}(\mathbf{t})\|$, which follow from (A.26) and (A.40), coincide. Thus, the estimates

$$(A.41) \quad \|\Psi_r^{\text{inv}}(\mathbf{t})\| \leq (r+1)^d \rho^{-r}, \quad r = 0, 1, 2, \dots, \quad \forall \mathbf{t} \in D(\mathbf{t}^0, \delta),$$

hold with the same ρ and d that in (A.32). Therefore, the series (A.33) converge locally uniformly in the disk $\{x : |x| < \rho_j(\mathbf{t})\}$ and represent the matrix function $\Psi^{\text{inv}}(x, \mathbf{t})$ which is holomorphic with respect to $x, \mathbf{t}$ for $x : |x| < \rho_j(\mathbf{t}^0), \mathbf{t} \in D(\mathbf{t}^0, \delta)$. The identities (A.34) for formal power series imply the identities

(A.42)

$$\Psi(x, \mathbf{t}) \Psi^{\text{inv}}(x, \mathbf{t}) = \Psi^{\text{inv}}(x, \mathbf{t}) \Psi(x, \mathbf{t}) = I \quad \forall x : |x| < \rho_j(\mathbf{t}^0), \mathbf{t} \in D(\mathbf{t}^0, \delta)$$

for the appropriate holomorphic matrix functions. Thus, the values of the matrix function $\Psi(x, \mathbf{t})$ for $x : |x| < \rho_j(\mathbf{t}^0), \mathbf{t} \in D(\mathbf{t}^0, \delta)$ are invertible, and $\Psi^{-1}(x, \mathbf{t}) = \Psi^{\text{inv}}(x, \mathbf{t})$.

Performing the change of variables $x \rightarrow x - t_j$, inverse to the change (A.2), we obtain the representation (6.3) for the local solution $Y_j(x, \mathbf{t})$, with $H_j(x, \mathbf{t}) = \Psi(x - t_j, \mathbf{t})$. The properties of the function $H_j(x, \mathbf{t})$, stated in the formulation of Proposition 6.3, express the properties of $\Psi(x, \mathbf{t})$, as established above. This completes the proof of Proposition 6.3. Q.E.D.

REFERENCES

- [AKNS] ABLOWITZ, M., KAUP, D., NEWELL, A. and H. SEGUR. *Nonlinear-evolution equations of physical significance*. Phys. Rev. Let., **32**:2, (1973), 125 - 127.
- [AnBo] ANOSOV, D.V. and A.A. BOLIBRUCH. *The Riemann - Hilbert Problem*. (Aspects of Mathematics, Vol. E 22), Vieweg, Braunschweig · Wiesbaden, 1994, i - ix + 190 pp.
- [Arn] АРНОЛЬД, В.И. *Кольцо когомологий группы крашенных кос*. Матем. заметки, 5:2, (1969) , с. 227 - 231. Engl. transl.: ARNOL'D, V.I. *The cohomology ring of the colored braid group*. Math. Notes, **5**, pp. 138 - 140 (1969).
- [Birk] BIRKHOFF, G. D. *The generalized Riemann problem for linear differential equations and the allied problems for linear difference and q-difference equations*. Proc. Amer. Acad. Arts and Sci., **49** (1913)
- [Bol1] БОЛИБРУХ, А. А. *Проблема Римана-Гильберта*. Успехи матем. наук, 45:2 (1990), с. 3 - 47 . Engl. transl.: BOLIBRUCH, A.A. *The Riemann-Hilbert problem*. Russian Math. Surveys, **45**:2 (1990), pp. 1 - 47.
- [Bol2] БОЛИБРУХ, А. А. *21-я проблема Гильберта для Фуксовых линейных систем*. Труды матем. института им. В.А. Стеклова, **206**. Москва, Наука, 1994., 160 с. Engl. transl.: BOLIBRUCH, A.A. *Hilbert's twenty first problem for Fuchsian systems*. Proc. Steklov Inst. Math, **5** (**206**), 1995, viii + 145 pp.
- [Bol3] БОЛИБРУХ, А. А. *Фуксовы Дифференциальные Уравнения и Голоморфные Расслоения*. (In Russian): BOLIBRUCH, A.A. *Fuchsian Differential Equations and Holomorphic Bundles*. МЦМО (Издательство Московского Центра непрерывного математического образования.), Moscow, 2000, 120 pp.
- [BoDoVe] BOUTET DE MONVEL, L., A.DOUADY and J.-L. VERDIER - editors. *Mathématique et Physique. Séminaire de l'Ecole Normale Supérieure 1979 - 1982*. (Progress in Mathematics, vol. **37**). Birkhäuser, Boston · Basel · Stuttgart, 1983
- [CoLe] CODDINGTON, E.A. and N. LEVINSON. *Theory of Ordinary Differential Equations*. McGraw Hill, New York·Toronto·London, 1955.
- [DIKZ] DEIFT, P., ITS, A., КАРАЕВ, А. and X. ZHOU : *On the Algebro-Geometric Integration of the Schlesinger Equations*. Commun. Math. Phys., **203** (1999), pp. 613 - 633.
- [Dick] DICKEY, L.A. *Soliton Equations and Hamiltonian Systems*. World Scientific. Singapore·New Jersey·London·Hong Kong, First Edition (Advanced Series in Mathematical Physics, Vol. 12) - 1991, ix+310 pp.; Second Edition (Advanced Series in Mathematical Physics, Vol. 26) - xii+420 pp., 2003.
- [DJKM1] DATE, E., M. JIMBO, M. KASHIWARA, and T. MIWA. *Transformation groups for soliton equations*. Proc. Japan. Acad. Ser. **A** Math. Sci.: I. **53**:1 (1977), pp. 6 - 10; II. **53**:5 (1977), pp. 147 - 152; III. **53**:5 (1977), pp. 153 - 158; IV. **53**:6 (1977), pp. 183 - 185; V. **53**:7 (1977), pp. 219 - 224; VI. **54**:1 (1978), pp. 1 - 5; VII. **54**:2 (1978), pp. 36 - 41.
- [DJKM2] DATE, E., M. JIMBO, M. KASHIWARA, and T. MIWA. *Solitons, τ functions and Euclidean Lie algebras*. Pp. 261 - 278 in: *Mathématique et Physique. Séminaire de l'Ecole Normale Supérieure 1979 - 1982*. BOUTET DE MONVEL, L., A. DOUADY, and J.-L. VERDIER - eds. (Progress in Math., **37**). Birkhäuser, Boston·Basel·Stuttgart, 1983
- [FuL] FUCHS, L. *Gesammelte Mathematische Werke. Band 3*. (Herausgegeben by R. FUCHS und L SCHLESINGER). Mayer & Müller, Berlin, 1909.
- [FuL1] FUCHS, L. *Zur Theorie der linearen Differentialgleichungen*. Sitzungsberichte der K. preuss. Akademie der Wissenschaften zu Berlin. Einleitung und No. 1 - 7, 1888, S. 1115 - 1126; No. 8 - 15, 1888, S.1273 - 1290; No. 16 - 21, 1889, S. 713 - 726; No. 22- 31, 1890, S. 21 - 38. Reprinted in: [FuL], S. 1 - 68.

- [FuL2] FUCHS, L. *Über lineare Differentialgleichungen, welche von Parametern unabhängige substitutionsgruppen besitzen.* Sitzungsberichte der K. preuss. Akademie der Wissenschaften zu Berlin. 1892, S. 157 - 176. Reprinted in: [FuL], S. 117 - 139.
- [FuL3] FUCHS, L. *Über lineare Differentialgleichungen, welche von Parametern unabhängige Substitutionsgruppen besitzen.* Sitzungsberichte der K. preuss. Akademie der Wissenschaften zu Berlin. Einleitung und No. 1 - 4, 1893, S. 975 - 988; No. 5 - 8, 1894, S. 1117 - 1127. Reprinted in: [FuL], S. 169 - 195.
- [FuL4] FUCHS, L. *Über die Abhängigkeit der Lösungen einer linearen differentialgleichung von den in den Coefficienten auftretenden Parametren.* Sitzungsberichte der K. preuss. Akademie der Wissenschaften zu Berlin. 1895, S. 905 - 920. Reprinted in: [FuL], S. 201 - 217.
- [FuR] FUCHS, R. *Sur quelques équations différentielles linéaires du second ordre.* Compt. Rend. de l'Académie des Sciences, Paris. **141** (1905), pp. 555 - 558.
- [Gant] ГАЙТМАХЕР, Ф. Р. *Теория матриц. 2-е издание.* Наука, 1966, 575 с. (In Russian). English transl.: GANTMACHER, F.R.. *THE THEORY OF MATRICES.* Chelsea, New York, 1959, 1960.
- [Gar] GARNIER, R. *Sur une classe d'équations différentielles dont les intégrales générales ont leurs points critiques fixes.* Compt. Rend. de l'Académie des Sciences, Paris. **151** (1910), pp. 205 - 208.
- [GKRM] GOHBERG, I., M.A. KAASHOEK, L. LEHRER and L. RODMAN. *Minimal divisors of rational matrix functions with prescribed zero and pole structures*, pp. 241 - 275 in: *Topics in Operator Theory, Systems and Networks.* (DYM, H. and I. GOHBERG- ed.) Operator Theory: Advances and Applications, **OT 12**, Birkhäuser, Basel Boston Stuttgart, 1984.
- [Ha] HARNAD, J., *Dual isomonodromic tau functions and determinants of integrable Fredholm operators.* In: *Random Matrix Models and Their Applications*, (Mathematical Sciences Research Institute Publications, **40**), BLEHER, P. and A. ITS - editors. Cambridge Univ. Press 2001.
- [HaI] HARNAD, J. and A. ITS. *Integrable Fredholm operators and dual isomonodromic deformations.* Commun. Math. Phys., **226** (2002), pp. 497 - 530.
- [Hart] HARTMAN, PH. *Ordinary Differential Equations.* Wiley, New York-London-Sydney, 1964. Russian transl.: ХАРТМАН, Ф. *Обыкновенные Дифференциальные уравнения.* Москва, Мир, 1970, 720 с.
- [HsSi] HSIEH, P., and Y. SIBIYA. *Basic Theory of Ordinary Differential Equations.* Springer-Verlag, New York-Berlin-Heidelberg, 1999, xi+468 pp.
- [Huk] HUKUHARA, M. *Ordinary Differential Equations.* (In Japanese). Iwanami-Zensho **116**, Iwanami-Shoten, 1950.
- [IKSY] IWASAKI, K., KIMURA, H., SHIMOMURA, SH. and M. YOSHIDA . *From Gauss to Painlevé. A Modern Theory of Special Functions.* (Aspects of mathematics: E , Vol. **16**). Vieweg, Braunschweig, 1991.
- [JMMS] JIMBO, M., MIWA, T., MÔRI, Y. and M. SATO. *Density matrix of an impenetrable Bose gas and the fifth Painlevé transcendent.* Physica **1D** (1980), pp. 80 - 158.
- [JMU] JIMBO, M., MIWA, T. and K. UENO. *Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. I. General theory and τ -function.* Physica **2D** (1981), pp. 306 - 352.
- [JM1] JIMBO, M. and T MIWA. *Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. II.* Physica **2D**, (1981), pp. 407 - 448.
- [JM2] JIMBO, M. and T MIWA. *Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. III.* Physica **4D**, (1981), pp. 26 - 46.

- [Kats1] KATSNELSON, V. *Fuchsian differential systems related to rational matrix functions in general position and the joint system realization*, pp. 117 - 143 in: Israel Mathematical Conference Proceedings, Vol. **11** (1997), *Proceedings of the Ashkelon Workshop on Complex Function Theory (May 1996)*, ZALCMAN, L. - editor.
- [Kats2] KATSNELSON, V. *Right and left joint system representation of a rational matrix function in general position (System representation theory for dummies)*, pp. 337 - 400 in: *Operator theory, system theory and related topics. (The Moshe Livšic anniversary volume. Proceedings of the Conference on Operator Theory held at Ben-Gurion University of the Negev, Beer-Sheva and in Rehovot, June 29 - July 4, 1997)*, ALPAY, D. and V. VINNIKOV - editors. (Operator Theory: Advances and Applications, vol. **123**), Birkhäuser Verlag, Basel, 2001.
- [KiKo] KITAIEV, A.V. and D.A. KOROTKIN. *On solution of the Schlesinger Equations in Terms of Θ -functions*. Intern. Math. Research Notes, 1998, No. **17**, pp. 877 - 905.
- [KoMa] КОРОТКИН, Д. А. и В. Б. МАТВЕЕВ. *О θ -функциональных решениях системы Шлезингера и уравнения Эрнста*. Функциональный анализ и его приложения, 34:4 (2000), с. 18 - 34 (Russian). English transl.: KOROTKIN, D.A. and V.B. MATVEEV. *On theta function solutions of the Schlesinger system and the Ernst equation*. Funk. Anal. Appl., **34:4**, (2000), pp. 252 - 264.
- [Mah] MAHOUX, G. *Introduction to the theory of isomonodromic deformations of linear ordinary differential equations with rational coefficients*. Pp. 35 - 76 in: *The Painlevé Property. One Century Later*. CONTE R. - editor. (CRM Series in Mathematical Physics.) Springer-Verlag, New York-Berlin-Heidelberg, 1999.
- [MSW] MASON, S.J., M.A. SINGER and N.M.J. WOODHOUSE. *Tau function and the twistor theory of integrable systems*. Journal of Geometry and Physics, **32** (2000), pp.397 - 430.
- [Miw] MIWA, T. *Painlevé property of monodromy preserving deformation equations and analyticity of τ functions*. Publ. RIMS Kyoto Univ., **17** (1981), pp. 703 - 721.
- [MJD] MIWA, T., M. JIMBO, and E. DATE. *Solitons. Differential equations, symmetries and infinite-dimensional algebras*. (Cambridge Tracts in Mathematics, **135**.) Cambridge University Press, Cambridge, 2000. x+108 pp.
- [New] NEWELL, A.C. *Solitons in Mathematics and Physics*. (CBMS-NSF Regional Conference Series in Applied Mathematics, **48**.) SIAM, Philadelphia, PA, 1985. xvi+244 pp. Russian transl.: НЬЮЭЛЛ, А. *Солитоны в Математике и Физике*. Мир, 1989, 326 сс.
- [Palm1] PALMER, J. *Determinants of Cauchy - Riemann operators as τ - functions*. Acta Applicandae Math, **18** (1990), pp. 199 - 223.
- [Palm2] PALMER, J. *Deformation analysis of matrix models*. Physica **98D** (1994), pp. 166 - 185.
- [Plem1] PLEMELJ, J. *Riemannsche Funktionenscharen mit gegebener Monodromiegruppe. (Riemann families with prescribed monodromy group)*. Monatshefte für Math. und Phys., **XIX**, (1908), pp. 211 - 245 .
- [Plem2] PLEMELJ, J. *Problems in the Sense of Riemann and Klein*. Interscience Publishers. A division of J. Wiley & Sons Inc., New York · London · Sidney, 1964, 175 pp.
- [Ra] RASCH, G. *Zur Theorie und Anwendung des Productintegrales*. Journ. für die reine und angew. Math., **171** (1934), pp.65 - 119.
- [Sakhn] САХНОВИЧ, Л.А. *О факторизации передаточной оператор-функции*. Доклады АН СССР, 226:4 (1976), с. 781 - 784. Engl. transl.: SAKHNOVICH, L.A., *On the factorization of an operator-valued transfer function*. Soviet. Math. Dokl. **17** (1976), pp. 203 - 207.

- [Sa] SATO, M. *The KP hierarchy and infinite-dimensional Grassman manifolds*. Proc. of Symposia in Pure Math., **49**:1 (1989), pp. 51 - 66.
- [SMJ1] SATO, M., MIWA, T and M. JIMBO. *Aspects of holonomic quantum fields. Isomonodromic deformations and Ising model*. Pp. 429 - 491 in *Complex Analysis, Microlocal Calculus and Relativistic Quantum Theory. Proceeding of the Colloquium held at Les Houches, Centre de Physique, September 1979*, IAGOL-NITZER, D. - editor. (Lectures Notes in Physics, Vol. **126**). Springer - Verlag, Berlin-Heidelberg-New York, 1980.
- [SMJ2] SATO, M., MIWA, T and M. JIMBO. *Holonomoc quantum fields. II*. Publ. RIMS Kyoto Univ. **15** (1979), pp. 201 - 278.
- [SaSa] SATO M. and Y. SATO. *Soliton equations as dynamical systems on infinite dimensional Grassmanin manifolds*. Pp. 259 - 271 in: *Nonlinear Partial Differential Equations in Applied Science; Proceedings of The U.S. - Japan Seminar, Tokyo, 1982*, (Lect. Notes in Num. Appl. Anal., **5**), 1982, FUJITA, H., P. LAX, and G. STRANG - eds.
- [Sch1] SCHLESINGER, L. *Über die Lösungen gewisser linearer Differentialgleichungen als Funktionen der singulären Punkte*. Journal für reine und angew. Math, **129** (1905), pp. 287 - 294.
- [Sch2] SCHLESINGER, L. *Vorlesungen über lineare Differentialgleichungen*. Leipzig und Berlin, 1908.
- [Sch3] SCHLESINGER, L. *Über eine Klasse von Differentialsystemen beliebiger Ordnung mit festen kritischen Punkten*. Journal für reine und angew. Math, **141** (1912), pp. 96 - 145.
- [SeWi] SEGAL, G. and G. WILSON. *Loop groups and equations of KdV type*. Publ. Mathem. IHES, № **61** (1985), pp. 5 - 65 . Reprinted in [TeUh], pp. 403 - 466. Russian transl.: СИГАЛ, Г. и ДЖ. ВИЛЬСОН. *Группа петель и уравнения типа КдФ*. Стр. 379—442 в книге: ПРЕССЛИ, Э. и Г. СИГАЛ. *Группы Петель*. Москва, Мир, 1990.
- [Sib] SIBUYA, Y. *Linear Differential Equations in the Complex Domain: Problems of Analytic Continuation*. (Translations of Math. Monographs, **82**), Amer. Math. Soc., Providence, Rhode Island, 1990, iv+267.
- [TeUh] TERNG, CH. L. and K. UHLENBECK - editors. *Surveys in Differential Geometry: Integral Systems. (Lectures in Geometry and Topology, sponsored by Lehigh University's Journal of Differential Geometry.)* International Press, Boston, 1998.
- [WNTB] WU, T.T., B. MCCOY, C. TRACY and E. BAROUCH. *Spin-spin correlation functions for the two-dimensional Ising model: Exact theory in the scaling region*. Phys. Rev. **B**, **13**:1 (1976), 316 - 374.
- [ZaSh] ЗАХАРОВ, В.Е. и А.Б. ШАБАТ. *Точная теория двумерной самофокусировки и одномерной модуляции волн в нелинейной среде*. Журн. Эксперим. Теор. Физ., 61:1, (1971), 118 - 134 (In Russian). English transl.: ZAKHAROV, V.E. and A.B. SHABAT (=A.B. ŠABAT). *Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in non-linear media*. Soviet Physics JETP, **34**:1 (1972), pp. 62 - 69.

DEPARTMENT OF MATHEMATICS, THE WEIZMANN INSTITUTE OF SCIENCE, REHOVOT, 76100, ISRAEL

E-mail address: `victor.katsnelson@weizmann.ac.il`

DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA, 84105, ISRAEL

E-mail address: `volok@math.bgu.ac.il`